

## RESEARCH ARTICLE

# A multi-method exploration of mindfulness as a coping tool: Perspectives from trauma-exposed, unhoused women residing at a drug treatment facility

Dana Rose Garfin<sup>1</sup> | Alexandra Amador<sup>2</sup> | Jessica Osorio<sup>3</sup> |  
Krista Sadiwa Ruivivar<sup>2</sup> | April Torres<sup>4</sup> | Adeline M. Nyamathi<sup>3</sup>

<sup>1</sup>Community Health Sciences, Fielding School of Public Health, University of California, Los Angeles, California, USA

<sup>2</sup>Department of Psychological Science, University of California, Irvine, California, USA

<sup>3</sup>Sue & Bill Gross School of Nursing, University of California, Irvine, California, USA

<sup>4</sup>HealthRIGHT360, Los Angeles, California, USA

## Correspondence

Dana Rose Garfin, Community Health Sciences, Fielding School of Public Health, 560 Charles E Young Drive South, Box 951772; Los Angeles, CA 90095, USA.  
Email: [dgarfin@ucla.edu](mailto:dgarfin@ucla.edu)

## Funding information

National Institute on Minority Health and Health Disparities, Grant/Award Number: MD013910

## Abstract

This multi-method study examined perspectives on mindfulness and coping strategies used by trauma-exposed women experiencing homelessness (WEH), residing in a state-funded residential drug treatment site in Southern California (United States). Questionnaires and in-depth focus group interviews were utilised to examine traumatic experiences over the lifespan, probable-posttraumatic stress disorder (PTSD), and coping strategies. Mindfulness was explored as a potential way to improve coping; potential benefits and challenges associated with implementing a mindfulness-based intervention (MBI) with trauma-exposed WEH were also investigated. A Community Advisory Board (CAB) was formed to identify key issues experienced by WEH and to develop a semi structured interview guide (SSIG). Using the SSIG, women participated in one of four focus groups (total  $N = 28$ ;  $n = 7$  per group). Quantitative data on demographic indicators, probable-PTSD, and trauma exposure were collected. Almost 90% of women met criteria for probable-PTSD; trauma exposure was exceedingly high; most women had experienced multiple traumas throughout their lives. Four main themes emerged from qualitative analyses, which drew from Grounded Theory and used *open*, *selective*, and *axial* coding: (1) ways of coping with trauma; (2) perspectives on mindfulness; (3) prior experiences with mindfulness; and (4) challenges for conducting a mindfulness programme. Overall, WEH used a variety of coping techniques to deal with their trauma, had some familiarity with mindfulness, and were optimistic an MBI would be helpful, despite identifying several challenges to implementation. MBIs may be helpful adjuncts to traditional care for trauma-exposed, WEH, recovering from substance use disorder. Population-specific considerations may improve implementation and participation.

## KEYWORDS

community-participatory research, coping, homelessness, mindfulness, PTSD, substance use disorder, trauma

## 1 | INTRODUCTION

Women experiencing homelessness (WEH) are at exceptionally high risk of trauma exposure, posttraumatic stress disorder (PTSD; North & Smith, 1992; Tsai et al., 2012), and co-occurring substance use disorder (SUD) (Guillén et al., 2020). Compared to the general population, among WEH, the prevalence of alcohol and drug use disorder is four and 12 times higher, respectively (Upshur et al., 2017). More generally, approximately 40% of those with SUD meet criteria for PTSD (Roberts et al., 2015). These comorbidities can be reinforcing: people may use substances as a means to 'forget' a trauma (Mills, 2009). Thus these comorbidities work in tandem to create additional risk factors for new traumas to occur (Duncan et al., 2019). Trauma may trigger substance use, and substance use may increase women's exposure to traumatic experiences. Therefore, the two issues are often mutually reinforcing, leading to a cycle of trauma exposure, PTSD, SUD, and high-risk situations that expose women to more trauma, exacerbating symptoms and resulting disparities.

Trauma and adverse childhood experiences (ACEs), commonly experienced in unhoused persons, are also associated with greater risk of homelessness (Duncan et al., 2019), chronic PTSD, and more severe SUD (Guillén et al., 2020), leading to social and occupational impairments that cause or maintain homelessness (Duncan et al., 2019). In prior assessments, 69% of WEH reported abuse as a child and 75% reported victimization as adults (Tsai et al., 2015); nearly 30% of unhoused individuals meet criteria for PTSD (Ayano et al., 2020). Moreover, the experience of homelessness itself is a traumatic experience (Phipps et al., 2019). WEH face a unique set of challenges (Phipps et al., 2019), which often include caring for children, avoiding current or former abuse perpetrators, and trying to maintain temporary or affordable permanent housing (Fonfield-Aylna, 2009). These challenges are reinforcing: for example, lack of childcare and/or regulations at temporary shelters can make interviewing for employment difficult for a women fleeing domestic abuse (Gültekin et al., 2014). Indeed, a large percentage of unhoused mothers have a history of social stressors and emotional and physical abuse, including a cycle of familial violence; they describe substance use as a way to cope with traumatic events, PTSD symptoms, and homelessness (Gültekin et al., 2014). Importantly, becoming housed does not always address problems including social integration, trauma, or SUD, which can in turn lead to future homelessness (Phipps et al., 2019).

Given such patterns and statistics, it is imperative WEH, particularly those with trauma exposure and associated PTSD and SUD, find effective coping skills that promote SUD recovery, help regulate PTSD symptomatology, and provide tools and resources for effective and sustained reintegration (e.g., obtaining employment and stable housing). Coping is defined as 'constantly changing cognitive and behavioural efforts to manage specific external and/or internal demands that are appraised as taxing or exceeding the resources of the person (pg. 141)' (Lazarus & Folkman, 1984). The Comprehensive Health Seeking and Coping Paradigm (CHSCP) provides a framework to alter, enhance and promote effective coping and service seeking

for those in need (Nyamathi, 1989; Nyamathi et al., 2021). In the CHSCP, coping is viewed as a transactional process, where social ecological factors (e.g., situational and personal factors, resources) influence an individual's cognitive appraisal, health goals, and in turn both immediate and long-term outcomes (Nyamathi, 1989). This improved coping is an essential component of reducing traumatic experiences such as violence (Boxer & Sloan-Power, 2013) that maintain PTSD symptoms and SUD. Understanding these processes, particularly from the perspective of the individuals seeking to improve their coping skills and adapt to challenging circumstances (e.g., PTSD, homelessness, SUD), is critical to supporting lasting recovery and behaviour change. Such information could help inform the design and administration of community-based services and interventions that target the development of improved coping skills, helping WEH to recover from PTSD, SUD, and homelessness.

A recent meta-synthesis of qualitative research on coping in persons experiencing homelessness with severe mental illness found that coping was a complex process (Karadzhev et al., 2020), noting that despite prolific research about coping with severe mental illness in general, the coping process in those living in disempowered conditions, including severe poverty and homelessness, has been understudied (Karadzhev et al., 2020). Moreover, the accumulation of adverse life experiences can thwart efforts to manage psychological comorbidities and ongoing life stressors (Padgett et al., 2012). These stressors may be particularly prominent during the reintegration process, where women face challenges including new employment, obtaining stable housing, learning more effective parenting skills, and ongoing legal issues.

### 1.1 | Coping in WEH with PTSD

Coping is a process, with dynamics that evolve according to life circumstances. Avoidant strategies focus attention away from stress-inducing stimuli or one's psychological/somatic reactions to it; non-avoidant strategies involve directing attention towards the source of stress (Suls & Fletcher, 1985). Classic meta-analytic findings support the efficacy of both types: some avoidant strategies are associated with short-term positive coping, while nonavoidant strategies tend to predict positive adaptation long term (Suls & Fletcher, 1985). In those with PTSD, avoidance, assessed using experimental designs (Hayes et al., 2004), and avoidant coping strategies and dissociation, assessed via self-report, were associated with increased PTSD symptomatology; self-report active-coping strategies were associated with resilience (Thompson et al., 2018). Relatedly, avoiding experiencing unpleasant events has been correlated with maladaptive psychological symptoms related to PTSD diagnoses (Hayes et al., 2004).

**Avoidant coping.** Some research on WEH suggests avoidant coping may be associated with poorer outcomes, including PTSD, SUD, and related psychological comorbidities. For example, prior research on previously homeless women with high ACEs found negative, avoidance-based coping (e.g., suppressing emotions, using substances) was common (Schmitt et al., 2021). Compared to housed

mothers, unhoused mothers reported higher stress scores, depression scores, and avoidant coping, with avoidant coping and depression (in turn comorbid with PTSD and substance use) positively correlated (Banyard & Graham-Bermann, 1998). In a study of 871 unhoused individuals, avoidant coping (e.g., withdrawing from others, sleeping more than usual) predicted mental distress (Nyamathi et al., 2000). Such data suggest that interventions that help survivors actively accept and address the experience of trauma and related symptoms may help reduce the stress response and promote sustained recovery. Of note, the complexity of stressors experienced in the context of multiple disadvantages suggests some avoidant coping strategies (e.g., avoiding temporary housing situations that may lead to re-traumatisation or feelings of helplessness or stigma) may, in fact, be adaptive (McMordie, 2021).

**Non-avoidant coping.** Approach- and acceptance-based coping strategies, including mindfulness-based approaches, may help alleviate the deleterious effects of trauma exposure and help break the cycle of repeated exposure to trauma, PTSD, and SUD. In a longitudinal sample of college students, increased approach-based coping was associated with decreased trauma exposure (Jenzer et al., 2020). In a sample of veterans with PTSD, positive cognitive coping was associated with decreased posttraumatic stress symptoms (Sheerin et al., 2018). Mindfulness, a cognitive process involving focussing one's attention on the present moment (Kabat-Zinn, 1990), has been conceptualised as a form of both approach and acceptance-based coping, and may help address the PTSD-SUD cycle. Theoretical reviews suggest mindfulness and acceptance-based strategies may be associated with resilience to PTSD and other psychological disorders including SUD (Witkiewitz et al., 2005). Accordingly, empirical data indicate that mindfulness mediates the relationship between PTSD and SUD severity (Bowen et al., 2017).

## 1.2 | Mindfulness-based interventions

Mindfulness-based interventions (MBIs) have been associated with positive outcomes in clinical samples with PTSD (Polusny et al., 2015) and SUD (Himelstein, 2011). In a sample of veterans with PTSD, compared to a control group, those who completed a Mindfulness-based Stress Reduction (MBSR) programme demonstrated increased mindfulness and decreased PTSD (Polusny et al., 2015). Relatedly, a scoping review reported moderate-large effect sizes for the relationship between MBIs and reduced PTSD (Boyd et al., 2018). In a study of adults in a SUD outpatient programme, Mindfulness-Based Relapse Prevention, compared to cognitive-behavioural relapse prevention and treatment as usual, resulted in reduced drug use and heavy drinking (Bowen et al., 2014). Such findings support the utility of implementing MBIs into treatment for those with co-occurring PTSD and SUD. Hence, community-based treatment centres could be the ideal environments to offer MBIs, as they may promote access, interest, and retention.

Although MBIs demonstrate promise for alleviating psychological maladies in unhoused individuals (Maddock et al., 2017), those

with SUD (Korecki et al., 2020), and other low-income populations experiencing high health disparities (Dutton et al., 2014), relatively little work has focussed on trauma, PTSD and SUD in WEH specifically. However, a recent systematic review on interventions to reduce PTSD and problematic substance use in survivors of interpersonal violence showed that teaching coping skills, including symptom stabilisation, emotional regulation, and social support, may be especially helpful for women with severe PTSD who have experienced chronic victimization (Bailey et al., 2019). Key components of MBIs include teaching skills that bolster emotion regulation, stress management, and leveraging group-based social support (Baer, 2003; Kabat-Zinn, 1990). Thus these interventions may be helpful at addressing the co-occurrence of PTSD and SUD in unhoused women, many of whom have experienced recurring victimization. This may help break the cycle of trauma and SUD that can perpetuate the cycle of homelessness. Moreover, components of MBI (e.g., community engagement, trigger awareness, increased control, self-acceptance) relevant to the *dual diagnosis* of PTSD and SUD may help facilitate an integrated *dual recovery* (Davidson et al., 2008). Yet, the extant literature has not explored mindfulness in the context of other coping techniques implemented by WEH, particularly using qualitative methods that can elucidate in-depth the experience of trauma, resilience, and recovery in this marginalised population.

## 1.3 | The present study

We explored coping techniques used by trauma-exposed WEH currently residing in a residential drug treatment site (RDTs). Using a community-based participatory approach (Israel et al., 2001), we used quantitative and qualitative data to describe traumatic experiences in WEH and coping techniques implemented during recovery from trauma, PTSD, homelessness, and SUD. Finally, we explored the use of mindfulness as a coping tool and the potential utility of implementing MBIs to promote recovery from PTSD and SUD in trauma-exposed WEH.

## 2 | METHOD

### 2.1 | Design

We utilised a multimethod design that implemented quantitative questionnaire data and in-depth focus groups (Maxwell et al., 2015). We used community-based research principles (Israel et al., 2001), informed by a participatory approach that included the use of a Community Advisory Board (CAB; Matthews et al., 2018). Quantitative questionnaire data ascertained descriptive statistics on the sample, including trauma exposure; qualitative data provided an exploration of key research questions regarding perspectives on mindfulness, MBIs and coping strategies related to trauma, SUD, and homelessness. All procedures were approved by the Institutional Review Board (IRB) at the University of California, Irvine.

## 2.2 | Setting, CAB sample, and procedure

Two RDTs in Southern California, funded under the same state umbrella organisation, participated in the study. The sites were comparable in terms of treatment administered (e.g., level of care, programs offered) and overall demographic composition; both had high diversity in client race/ethnicity and age. RDTs were funded primarily by Drug Medi-Cal, a treatment funding mechanism for eligible Medi-Cal members whose goal is to 'provide eligible Medi-Cal members with access to the care and services they need for a sustainable and successful recovery' from SUD (California Department of Health Care Services, 2022). Average stay length at the RDTs typically ranged from 60 to 120 days, with 24–25 h per week of Drug-Medi-Cal treatment, a combination of evidence based clinical practice, 'soft skills,' and individual sessions with therapists, case managers, and SUD counsellors. Formal MBIs were not available, although mindfulness-related skills were offered as part of informal client education at one of the RDTs.

CABs consist of community members who provide consultations to improve capacity and quality of research projects (Matthews et al., 2018). Our CAB included (a) six trauma-exposed WEH, living in a RDTs, (b) one substance abuse counsellor, (c) a site licenced clinical psychologist, (d) a site clinical manager, and (e) a site clinical director. CAB members were drawn equally from both sites; two meetings were conducted over Zoom conferencing with research team members present at both sites; the principal investigator (PI) alternated onsite presence.

## 2.3 | Semi-structured interview guide (SSIG)

Together, the CAB and research team designed a SSIG that guided subsequent focus groups. CAB members were compensated \$10 per session. A key goal was to obtain WEH's perspective on MBSR, which typically has an 8-week format, including a 1-day retreat (Kabat-Zinn, 1990). MBSR was selected as it is one of the most successful and widely implemented MBIs, safely and effectively implemented in populations with PTSD (Polusny et al., 2015), and served as the foundation for a variety of subsequent interventions used to treat clinical conditions relevant to trauma-exposed WEH with SUD (e.g., Acceptance and Commitment Therapy, Dialectical Behavior Therapy, Mindfulness-Based Relapse Prevention, Mindfulness Oriented Recovery Enhancement) (Korecki et al., 2020). MBSR involves completing homework assignments including guided meditations. Key practices include yoga, body scan practice, and breath meditation.

Questions from the SSIG included: What are your prior experiences with mindfulness activities such as yoga, body awareness, and guided meditations? What coping skills have you used in the past to deal with your trauma? The full, finalised SSIG is included in Supplement 1.

Using IRB-approved flyers, RDTs clinical staff informed residents of the opportunity to participate in focus groups. The research team screened interested women. Eligibility included: age over 18,

experienced homelessness in the last six months, and exposed to a DSM-5 criterion A trauma (American Psychiatric Association, 2013).

## 2.4 | Focus group sample, setting and procedure

In total, 35 women were screened; 32 were eligible. A convenience sample of 28 WEH participated in the focus groups ( $N = 4$ ). Participants in the focus group were distinct from CAB members to differentiate between the research design process (a function of the CAB) and data collection (which occurred during the focus groups) to minimise potential bias. Of eligible WEH who did not participate, one had a court case and two had conflicting off site appointments; one decided not to participate. Setting is the same as above. Flyers were distributed throughout the RDTs by site staff, informing women of the opportunity to participate.

Focus groups were selected as a methodology over individual interviews as randomized, experimental work shows that focus groups can more effectively facilitate disclosure of sensitive and personal information (Guest et al., 2017). They may also be better for exploring clinical patients' perspectives (Coenen et al., 2012) as participants may feel more supported and less stigmatised when sharing in a group. Indeed, in some populations, the group-based context of the focus group may enable greater comfort and reassurance and a feeling that one's difficult experiences are shared by others; in those instances, participants may find focus group discussions a positive experience (Jordan et al., 2007). Focus groups can also provide a safe space for vulnerable women to share their experiences without fear of criticism, can maximise the perspectives of individuals, and can empower individuals of the group to speak, rather than focus on responding to the researcher's question in a one-one-one interview (Owen, 2001). We took many steps to ensure the emotional safety of participants: everyone was reminded to keep what was shared confidential, pseudonyms were used, site staff was available at any time should participants need further psychological assistance, and participants were informed their participation was voluntary and they could stop at any time.

Using the SSIG, focus groups, each with seven WEH, were led by the PI at a private location at the RDTs. MBSR activities were demonstrated using a theatre-style approach; WEH did not complete the MBSR course but were asked to comment on the acceptability of its key components. Focus groups were audio recorded; two assistant moderators took detailed notes. Pseudonyms were created to protect anonymity. A trained therapist was available onsite if any adverse events or extreme distress occurred. Focus groups lasted 1.5–2 h. Women were compensated \$15.

## 2.5 | Quantitative measures

Participants completed a sociodemographic questionnaire (see Supplement 2) and the *Life Events Checklist* (Gray et al., 2004), a measure of adverse life experience with good psychometric properties.

Participants were asked when in the lifespan each event occurred (i.e., before age 18, 18 or older, or both). They also completed the *Primary Care PTSD Screen (PC-PTSD)*, a five-item screening tool that has been validated against clinician diagnostic interviews (Prins et al., 2015, 2016).

## 2.6 | Qualitative analyses

Focus groups were professionally transcribed; the PI and three research assistants (RAs) involved in the focus groups met to engage in precoding activities. First, each RA independently reviewed four transcripts (one from each focus group) in entirety. Key constructs were highlighted using *open coding* techniques, with an inductive research process that drew from Grounded Theory (Glaser & Strauss, 1967), whereby the codes themselves emerged from the data (Glaser & Strauss, 1967; Linneberg & Korsgaard, 2019; Williams & Moser, 2019).

*Open coding* (Williams & Moser, 2019) was used to identify broad and distinct concepts using the full transcripts from the focus group, derived from words, phrases and 'in vivo' coding (Manning, 2017), where an exact phrase from a participant's response was used to define a code. After this initial list of codes was generated, using *axial coding*, a process that examines open codes to find those most interrelation or overlap, transcripts were re-reviewed and themes were refined, aligned and categorised (Williams & Moser, 2019). Then, using *selective coding*, a process where categories of data organised during axial coding are combined conceptually and expanded thematically, data were organised into cohesive and meaning-filled expressions (Williams & Moser, 2019). During the *selective coding* process, quotes (and associated codes) were placed in a unique tab in Excel, denoting the potential theme. Subthemes were derived from the axial coding process. Axial and selective coding procedures used a hybrid model of inductive and deductive processes (Linneberg & Korsgaard, 2019), where concepts derived from the SSIG, CAB discussions, and stress and coping theory (Lazarus & Folkman, 1984; Suls & Fletcher, 1985), as applicable, were used to inform emerging themes and subthemes.

For example, one quote from the focus group was, 'My drug use that I was coping with had got out of hand and I ended up being in the psych ward.' During open coding, this was defined as 'using drugs and alcohol to cope' and 'self-medicating.' We also noted the participant described this method as 'ineffective.' Next, during the axial coding process, this code/quote was grouped into 'avoidant coping,' which also became a subtheme. During the selective coding process, 'avoidant coping,' was grouped with other coping techniques into one of the main themes: 'ways of coping with trauma.'

RAs used written notes and audio recordings from the focus groups for further contextualisation. The RAs and the Principal Investigator (PI) met at least weekly during the coding processes to review transcripts line-by-line discuss potential codes and themes. At each stage in the open, selective, and axial coding process, RAs coded individually and then discussed their codes with the group. Through this iterative process, detailed meeting notes were kept and coding

disagreements were resolved by all three coders and the PI via discussion and consensus (Campbell et al., 2013).

## 3 | RESULTS

### 3.1 | The sample

Participants were, on average 30.29 years old ( $SD = 8.6$ , range = 18–59) and reported diverse racial/ethnic identities: White ( $n = 10$ , 35.7%), Black ( $n = 5$ , 17.9%), and Hispanic ( $n = 13$ , 46.4%); two (7.1%) additionally identified as part Native American. Educational attainment was relatively low, although variability existed: less than high school ( $n = 11$ , 39.3%), high school/GED ( $n = 6$ , 21.4%), some college ( $n = 10$ , 35.7%), and bachelor's degree ( $n = 1$ , 3.6%). One (3.6%) WEH who completed high school/GED also reported vocational training.

All 28 WEH had been diagnosed with an SUD and had been involved with the justice system, with 22 (78.6%) also having received a doctor diagnosed mental health condition. These included depression ( $n = 18$ , 64.3%), anxiety ( $n = 18$ , 64.3%), PTSD ( $n = 16$ , 57.1%), bipolar disorder ( $n = 4$ , 14.3%), attention deficit disorder ( $n = 3$ , 10.7%), schizophrenia ( $n = 4$ , 14.3%), borderline personality disorder ( $n = 4$ , 14.3%), and obsessive compulsive disorder ( $n = 1$ , 3.5%). Of the participants, 17 (60.7%) identified as heterosexual, 4 (14.3%) identified as lesbian, 6 (21.4%) as bisexual, and one 1 refused. Most of the women had children ( $n = 25$ , 89.3%), with one woman currently pregnant.

### 3.2 | Quantitative results

#### 3.2.1 | Trauma exposure and prevalence of PTSD

Table 1 presents trauma exposure prevalence. Exposure was exceedingly high both in childhood and adulthood. Women were also provided the opportunity to report traumatic events not specifically assessed in the trauma inventory. Those reported included: cracked skull, held at gunpoint, miscarriage, and parental conflict. Three women specified losing their children to Child Protective Services, three specified domestic violence, and one indicated extreme stress about 'being gay.' Of participants, 89% met DSM-5 criteria for probable PTSD.

### 3.3 | Qualitative results

#### 3.3.1 | Overview

During the initial open coding procedure, 212 potential codes were generated. During axial coding, 38 codes were identified, and then pared down into eight themes. Four of these themes related to the coping process including perspectives and experiences with mindfulness and are presented in this manuscript. (The additional four

TABLE 1 Exposure to negative life events across the lifespan (N=28)

Event	Lifetime		Childhood before age 18		Adulthood, age 18 or higher	
	%	n	%	n	%	n
Physical assault	96.0	27	53.6	15	82.1	23
Sexual assault (rape, attempted rape, made to perform another sexual act)	85.0	24	53.6	15	67.9	19
Assault with a weapon	82.1	23	39.3	11	64.3	18
Other [besides sexual assault] unwanted sexual experience	71.4	20	42.9	12	57.1	16
Transportation accident	67.9	19	42.9	12	50.0	14
Captivity (kidnapped, abducted, POW, etc)	57.1	16	14.3	4	50.0	14
Serious accident at work, home or during recreation activity	53.6	15	32.1	9	25.0	7
Sudden violent death	46.4	13	21.4	6	35.7	10
Severe human suffering	42.9	12	25.0	7	32.1	9
Sudden accidental death	39.3	11	10.7	3	32.1	9
Fire or explosion	35.7	10	21.4	6	14.3	4
Serious injury, harm or death you caused to other	32.1	9	7.1	2	28.6	8
Life-threatening illness or injury	28.6	8	3.6	1	25.0	7
Exposure to toxic substance	17.9	5	3.57	1	17.9	5
Combat	14.3	4	0.0	0	14.3	4
Any other very stressful event or experience	64.3	18	32.1	9	53.6	15

themes centred specifically on the perceived acceptability and feasibility of a mindfulness programme, are conceptually distinct, and beyond the scope of the present analyses.) The four themes were: (1) ways of coping with trauma; (2) perspectives on mindfulness; (3) prior experiences with mindfulness, and (4) challenges for conducting a mindfulness programme.

#### **Theme 1. Ways of Coping with Trauma.**

Women described a variety of strategies for coping with their trauma, relevant to the experience of homelessness, trauma, and the recovery process. These coping strategies included those used prior to, during, and subsequent to the experience of homelessness. These strategies were grouped into the subthemes of (1) avoidant coping; (2) emotion-focussed coping; (3) proactive coping; and (4) social support seeking.

**Subtheme 1: Avoidant Coping.** Avoidant coping techniques were relevant to chronic and repeated experiences of homelessness, trauma, and difficulty with the recovery process. These included eating, sleeping, using drugs and alcohol, having sex, distraction (e.g., playing video games) and isolating from others. The WEH mostly described avoidant techniques as 'ineffective' for managing trauma and stress and as commonly used prior to coming to treatment.

"I was self-medicating on drugs. That was my biggest coping skill. ...I came into [treatment facility] hoping to learn new coping skills because I was self-medicating. So, drugs was my coping skills for like three years."- WEH1

"I would hide from everything. I would just run from it and just going keep my mind occupied with other things that didn't matter. Like, you know, like going and using or just skipping, you know, the reality of what just happened. Like, I would just hide from it. ...." - WEH2

"Well, I used drugs in the past so that was, but that was a coping skill. But I'm learning healthy coping skills now, but that was when I using." WEH3

"Not just that but like it was, it was ineffective, and like it only built on to like, the shame or guilt that you were already feeling. Like, for like in my example. Usually, I don't get so much angry as I do get ashamed. So, hiding away from it only builds on like, the stress of it, because I haven't faced whatever, whatever it is. So, it gets stronger, and the fact that I would do drugs, or try to run away from my problems, or drink whatever." WEH4

**Subtheme 2: Emotion-focussed Coping.** Emotion-focussed coping was commonly implemented in response to experiencing a stressor or a stressful emotion and included mainly healthy activities. Examples included self-care (e.g., taking a hot bath or shower), reading, writing, expressing emotions, prayer, journaling, listening to music, reading, gardening, self-reflection, exercise, breathing, or

talking to a therapist. The use of art was also frequently reported. Some WEH reported 'expressing emotions when they come up' or allowing themselves to cry.

"I write out stuff to people and talk about it to get it off my chest instead of holding it in because I have a bad problem doing that." -WEH4

"For me, like, I was very, like quick to react to things and angry and like, I was a little out of control. But being in [treatment facility], ...they taught me like the breathing and calming down because I have really bad anxiety. -- And I practice it and I never thought that it worked before. But I know if you really practice it and want to calm down and change and you do what they say and follow direction and breathe and do the, you know, just breathe, breathing...helps me... It calms me down a lot." -WEH5

"Expressing the emotions when they come up. Don't hold them in. So, I just have like a couple people around here that I go to, and I voice and vent to." - WEH1

**Subtheme 3: Proactive Coping.** Six women mentioned proactive coping techniques including avoiding negative influences, exploring healthy alternatives to negative or avoidant coping (e.g., using a nicotine patch to quit smoking), and practicing new coping skills taught in treatment. These coping skills related to techniques that could be done *in advance* of experiencing stress in order to facilitate a better response to it.

"Focus, setting your intention. For us, I mean, we have a lot of trauma. We have a lot of PTSD. We have a lot of flashbacks, bad dreams. We have all of those, those symptoms of our trauma. And so, when you become aware of your body and your mind and where it's taking you, you can control it. And when you've learnt to practice that control, practice makes perfect. So, you become a perfectionist in controlling your mind to not detour into those thoughts." -WEH6

"You know, I just accept what happened to me. When I came to this programme, I accepted a lot of things. And so, with accepting ...[what] happened to me and that... wasn't okay, now I move forward. It's okay for me because I can't just keep on thinking... now I know not to do things or be in certain places." -WEH2

**Subtheme 4: Seeking Social Support.** Learning how to effectively utilise social support was often discussed as a new skill learnt during treatment to help with traumatic stress symptoms and memories, drug and alcohol cravings, and other sources of general stress and distress. Main sources of social support were peers at the RDTS,

family (specifically mothers and sisters), and therapists; social support was often used in conjunction with emotion-focussed coping.

"I usually talk to my mom or to my sister or here to my therapist, you know, and I let her know how I'm feeling when it comes to that."-WEH5

"But one of like the new coping skills I'm using is like talking to my peers. Because normally, I don't like talking to people because I don't want to be a burden on them. But just knowing that I have somebody that will listen to me as like, I talked to them and sometimes they don't even say anything back. It just makes me feel like, like I just got a whole lot of stuff off my chest." -WEH6

### **Theme 2. Perspectives on Mindfulness.**

While many women were familiar with the term 'mindfulness,' they had various conceptualisation of its definition and use. These subthemes were: (1) mindfulness as a calming technique; (2) mindfulness as a grounding technique; (3) non-judgement and self-acceptance; and (4) respect and awareness for others and surroundings.

**Subtheme 1: Mindfulness as A Calming Technique.** Some participants were familiar with the term mindfulness and related techniques. This primarily related to using mindfulness as a component of an emotion regulation strategy to promote stress reduction and less reactivity, decrease experiences of anger and aggression, and enhance feelings of calm.

"I feel like it kind of just, it helps you like soothe your mind. So, instead of you like getting to a point where if you're angry and then just going off on somebody and exploding, you can take that and just meditate on it and calm down and be able to walk away and then it can give you like a piece of mind, kind of a thing."-WEH7

"It [meditation] relaxes me a lot. Like, especially if I'm like in a heated moment, to meditate and just collect myself and get a hold of like, me, it helps." -WEH5

"So, I really like the use of like mindfulness techniques like meditation and yoga.....they allow your body to relax and release tense emotions, and you might feel heartbreak throughout the day. And it, and it becomes easier, especially during the nighttime when you find it hard to sleep because you find your mind like, like, racing. So, utilising mindful techniques I feel, allows you to release the stress of the day. Kind of put your name on it and be done with it." -WEH4

**Subtheme 2: Mindfulness as a Grounding Technique.** Relatedly, WEH talked about using mindfulness as a tool to reorient the mind

towards present-moment awareness and find psychological safety during times of stress. One WEH specifically stated, 'I feel like it's [mindfulness is] a grounding technique' while another stated 'they [mindfulness techniques] put you in a state of what you, where you, are now.'

"I think mindfulness, the first thing that comes to me is to bring myself safe." -WEH8

"And if your mind's too, like, going at the same time like, you're thinking about too much, you can come back to like what's going on and use your five senses and kind of just not worry so much about what happened or what's going on. Just worry about that, like she said, present moment." -WEH9

**Subtheme 3: Non-judgement and Self-acceptance.** Several women mentioned mindfulness being related to non-judgement of self or feelings, and that process being a tool to 'feel better' during times of stress.

"Being mindfulness is like, having no judgement on your feelings as you feel. It's kind of like you're accepting what you're feeling and that moment so you're aware of that. You're not going to judge that. You can go through the process and then eventually, you'll feel better." -WEH10

"Yeah, you don't judge your own judging." -WEH4

**Subtheme 4: Respect and Awareness for Others and Surroundings.** Women often described mindfulness as a component of being respectful for the people around them and their surroundings more generally. Some described being mindful as 'thinking of others.'

"To be in the now, to know what's going on at that moment ...so you're aware of your surroundings, you're aware of what you're doing." -WEH9

"Yeah, you're like aware of your feelings, your emotions, the smell, the flowers, the trees. ...you're also mindful of, you know, ...respecting, okay, they're sleeping, I'm going to be mindful and not be loud. To be generous." -WEH5

### **Theme 3. Prior Experiences with Mindfulness.**

Many of the women reported prior experiences with mindfulness and related practices. The subthemes were: 1) familiarity with mindfulness as a technique; 2) mindfulness as a coping technique; 3) mindfulness for relaxation; and 4) mindfulness as a component of faith practice.

**Subtheme 1: Familiarity with Mindfulness as a Technique.** Many of the women expressed familiarity with mindfulness in the past,

including learning about it while incarcerated or in treatment. Many of them had heard of or participated in sitting meditation, yoga, daily reflections, and body awareness activities.

"Because when I was in jail, I went to like the meditation class or whatever. And that's what the teacher said to do. She said start off small like two minutes." -WEH8

"In some of our classes, we do it. Like, some of the, like, facilitators will read something...but like I haven't had a whole lot. I've been to like a couple of AA meetings that were meditation meetings and those are pretty cool because they like read from a little book. Like, you meditate and then you read from the little book and you meditate again and then you talk about it and you meditate again." -WEH12

**Subtheme 2: Mindfulness as a Coping Technique.** The women described mindfulness as helpful for coping with stress and other maladies (e.g., insomnia, physical pain) and downregulating during times of distress (e.g., feeling upset).

"It helped me deal with like, physical health issues as well as mental." -WEH1

"I used to ...go to sleep when I couldn't fall asleep. I used to do the body scan." -WEH14

"And that meditation, ...I like breathe and then I try to like clear my head. And when I clear my head, I'm just like, clear my head and then I come down. Like, for me being like angry or wanting to like hurt them, it just like goes away. It goes away. So, without even realizing that it was helping me, it was helping me." -WEH15

**Subtheme 3: Mindfulness for Relaxation.** Relatedly, some reported familiarity with mindfulness as a tool for relaxation during times of stress, for anxiety reduction, and to not 'keep things bottled up.' They described their use of mindfulness-related practices during times of distress and as a way to experience and process difficult emotions.

"I used it as an anxiety releaser." -WEH16

"So, I really like the use of like mindfulness techniques, like meditation and yoga. Not just because they put you in your state of ...where you are now, but they allow your body to relax and release tense emotions, and you might feel heartbreak throughout the day. And it, and it becomes easier, especially during the night-time when you find it hard to sleep because you find your mind like, like, racing. So, utilising mindful



techniques I feel, allows you to release the stress of the day. Kind of put your name on it and be done with it." - WEH17

#### **Subtheme 4: Mindfulness as a Component of Faith Practice.**

Several of the women discussed mindfulness in the context of their faith and other spiritual practices. More specifically, they reported exposures to the concept of mindfulness (mostly meditation practices) that were incorporated into Church or other faith-related settings.

"I go to the faith-based meditation that they have here and listen to church type music and it's very relaxing." - WEH1

"I also go to the faith-based [meditation], and it's also relaxing, and it helps me a lot." -WEH18

#### **Theme 4. Challenges for Conducting a Mindfulness Programme.**

Although overall the women were positive about MBIs, they noted potential challenges to implementation and provided general feedback regarding conducting MBSR with this population. These subthemes were: (1) difficulty focussing; (2) difficulty trusting others; (3) self-judgement; (4) trigger awareness; and (5) challenges to participation.

**Subtheme 1: Difficulty Focussing.** Many WEH reported that, based on their experiences during the demonstration and during prior experiences with mindfulness practices, difficulty focussing during the MBSR activities, including mediation practices and class discussions, may be an issue due to internal cognitive distractions or environmental noise. Other difficulties included sitting in 'one place too long' and that in treatment sometimes 'it's just hard to focus.'

"I've never done that before. I tried to do that, but I can't focus or even concentrate on doing it. It's one of my big things. Like, I can't concentrate. I can't focus on it." -WEH11

"So, I really want to learn how to, you know, meditate and whatnot. Because the only thing I know how to do right now is just breathe or take a walk or cry because I can't get into the yoga thing. I can't focus. I cannot focus. And then like, when there's a noise like, it's bad. It just got me out of it, you know. So, for me that's really hard." -WEH19

**Subtheme 2: Difficulty Trusting Others.** Many WEH reported that their potential difficulty trusting others should be considered when conducting group-based interventions. Key concerns involved not identifying with other RDTs residents, low trust from living on the streets, and feeling others did not care about their experience or perspective.

"Sometimes I feel like even talking to them [other residents] is pointless. So, I'd just rather just shut off ... I'm not going to say nothing because I know it's not going to mean nothing." -WEH2

**Subtheme 3: Self-judgement.** Another concern related to the high level of global and specific self-judgement many of the women reported. Such concerns were relevant to negative self-oriented cognitions, forgetting to praise oneself, or feeling stupid. Self-judgement was described with respect to trying a new activity such as yoga, difficulty focussing during instruction, and feeling like their prior drug use caused cognitive impairments.

"Like, we're not able to focus on that because we've been through so much. And it's like, hard for us get our mental like, right. I feel like even though I've been here for like almost three months...I can understand what they're saying." -WEH9

**Subtheme 4: Trigger Awareness.** Many WEH emphasised the importance of being aware of triggers, including memories or references to prior trauma exposure and substance use, and issues currently being unpacked as part of individual psychotherapy. For example, one WEH referenced water bottles and pens as examples of everyday objects formerly used as drug paraphernalia. One veteran referenced being outside as a potential trigger as it reminded her of her prior combat experience. Another WEH noted many in treatment were reconciling 'harsh' matters from the past and were quite raw emotionally. Simultaneously, some women expressed optimism that mindfulness and related techniques could be useful tools to face and overcome triggers, although creating a 'safe space' was critical to that effort.

"If you're drug addict, it's every little thing can become triggers." -WEH20

"I don't like closing my eyes because if I close my eyes, I get flashbacks. I see everything again. I don't feel comfortable with that." -WEH21

**Subtheme 5: Challenges to Participation.** Despite overall positive feedback, challenges to conducting MBSR with trauma exposed WEH were identified. A key concern was inspiring participants to enrol in the programme and engage in the practices. Many suggested a reward for participation but cautioned that could also lead to recruiting people who are not serious about the programme. Other concerns included meditation length (e.g., 45-min meditations might be too long) and scheduling MBSR classes without conflicting with other RDTs requirements.

"Because, you know, either you're going to take it seriously or you're not. In those who take it serious will

probably come back and those who don't, probably won't come back the next time, you know." – WEH14

## 4 | DISCUSSION

Herein, in a sample of WEH living in a residential drug treatment programme, we documented the experience of trauma throughout the lifespan. We explored coping skills implemented in response to that trauma and used qualitative methods to elucidate WEH's perceptions about the benefits and detriments of such skills. On balance, approach, proactive, and emotion-focussed coping seemed to help promote adaptive behaviours; avoidant strategies were generally discussed in the context of maladaptive strategies for dealing with homelessness, trauma, and SUD. Moreover, we investigated the potential of MBIs to aid in recovery from addiction and trauma in WEH, finding that mindfulness-based practices were viewed positively, and MBIs received enthusiastically despite several challenges that were identified. By using both quantitative and qualitative methods, we provide a rich context for understanding the process of recovery from trauma and related stressors in trauma-exposed, WEH. Such a multi-method approach helps inform 'participatory action research,' by providing an integrative approach to understanding and addressing complex social issues (Ivankova & Wingo, 2018), including the intersection of and recovery from trauma, homelessness, and substance abuse.

Quantitative data illustrated the exceedingly high level of trauma exposure experienced by WEH throughout the lifespan. Strikingly, over 96% reported physical assault and over 85% reported sexual assault; nearly 60% reported being held in captivity. Traumas occurred early in life (before age 18) and in adulthood, in alignment with prior research documenting high trauma exposure throughout the lifespan in those experiencing homelessness, leading to associated negative psychiatric outcomes (Tsai et al., 2015). Research on unhoused parent-child dyads demonstrates intergenerational relationships between parent-childhood adversity, parent-adulthood adversity, child adversity, and child-traumatic symptomatology (Lafavor et al., 2020). This intergenerational cycle is reinforced through both environmental and biological/genetic processes that maintain PTSD symptoms (Yehuda et al., 2016), perpetuating health disparities.

As with prior research (Karadzov et al., 2020), our data revealed WEH engaged in a range of coping behaviours to manage stressful events and PTSD, SUD, and other mental health ailments. Women tended to discuss avoidant coping in the context of unhealthy behaviour patterns (e.g. numbing with drugs, alcohol, or sex), similar to research suggesting avoidant strategies may be associated with deleterious outcomes and PTSD symptom maintenance (Thompson et al., 2018). Since avoidant coping involves actions and thoughts that evade confrontation with stressful events (Wu et al., 2013), such coping could thwart efforts in treatment aimed at teaching healthier coping skills. In contrast, approach-based coping strategies and those focussing on emotion-regulation and social support were discussed by WEH as facilitating recovery, psychological adjustment, and

healing. This aligns with research on trauma-exposed patients finding active coping strategies positively correlate with resilience and negatively correlate with PTSD (Thompson et al., 2018). Using art as a coping tool was discussed positively by the participants, similar to work suggesting art-based therapies can enhance resilience and reduce PTSD (Schouten et al., 2019).

WEH generally reported familiarity with the term mindfulness and related practices (e.g., yoga, sitting meditation), describing mindfulness as proactive or emotion-focussed coping, in alignment with work suggesting mindfulness practices can be viewed in the context of approach and acceptance-based coping (Thompson et al., 2011). Overall, mindfulness practices were viewed positively, in accordance with research indicating feasibility and acceptability of MBIs with homeless adults (Maddock et al., 2017) and populations experiencing high health disparities more generally (Dutton et al., 2014). Several challenges to implementation were identified, related to both mindfulness practices (e.g., difficulty focussing) and administering an intervention at an RDTS (e.g., difficulty trusting others, trigger awareness). Suggestions for overcoming barriers were noted (e.g., incorporate rewards, shorten practices, onsite therapist availability). Such findings provide useful data for designing and implementing MBIs with trauma-exposed WEH.

Participants reported that engaging in mindfulness-related practices (such as yoga, breathing, meditation, and reflection) were helpful for downregulating heightened emotional states including anger, anxiety, and stress. Such self-report data aligns with prior empirical work suggesting that MBIs help improve biological indicators of the stress-response (Bower et al., 2017) and may help lower physiological arousal (Lang et al., 2012). Such processes are critical for reducing traumatic-stress-related symptomatology, as data suggests biological dysregulation may precede the development of – and at a minimum maintain – PTSD symptoms (van Zuiden et al., 2013). Relatedly, research suggests mindfulness and acceptance are associated with post-trauma resilience (Thompson et al., 2011). This provides evidence that offering MBIs and related practices in treatment facilities serving women experiencing high health disparities could encourage a broader range of coping strategies to help deal with the stressors of recovery and re-integration (e.g., transition to sober living, new employment, parenting challenges). It also bolsters research with homeless youth (Sibinga et al., 2011), unhoused adults (Maddock et al., 2017) as well as WEH and young children (Alhusen et al., 2017) showing that MBIs could help with downregulation of difficult emotions, facilitate better coping, and improve mental health.

Such tools may also be concurrently helpful at alleviating co-occurring SUD, particularly with trauma-exposed WEH. The concept of integrating mindfulness practices into SUD treatment has existed for decades, given that cognitive and behavioural processes associated with mindfulness are integral to promoting abstinence and reducing relapse (Breslin et al., 2002). These processes include increased confidence for managing stressors, awareness of relapse triggers and decreased drug-cue relapse, dealing with negative affect, increased confidence, and acceptance of the past with commitment

to change (Breslin et al., 2002). Lack of effective tools for coping with interpersonal stress and negative affect contributes to relapses (Breslin et al., 2002), evidenced in daily diary studies (Ayer et al., 2011) and epidemiological work (Keyes et al., 2011). Such experiences were reported by the women in our focus groups. From a mechanistic perspective, mindfulness may increase awareness of potential triggers (Breslin et al., 2002), contributing to improved monitoring of situations that may promote relapse and destination to other relapse triggers over time. Resultingly, data suggests that mindfulness-based relapse prevention strategies that teach stress management, cue and trigger awareness, and effective coping skills to regulate negative affect and reactivity may be helpful to promote abstinence and prevent relapse (Bowen et al., 2014; Witkiewitz et al., 2005). Hence, in the context of the findings presented herein, MBIs may be useful to integrate into services for unhoused individuals.

However, such MBI programs may be best administered at an RDTS or other homeless service setting that could consistently administer the intervention over time, as prior research has demonstrated the importance of ongoing provider support and comprehensive and integrated services when administering trauma-informed care to those experiencing homelessness (Hopper et al., 2010). Moreover, based on participant's feedback, MBI practices readily lend themselves to a gender-responsive treatment model (Saxena et al., 2014). For example, the intervention can be designed (including group size, trigger awareness, trainer characteristics) to account for the specific needs of trauma-exposed women (e.g., victimization, motherhood) in the context of the gender-specific stress and stigma (e.g., felt invisibility) of experienced homelessness for women (Whitzman, 2006).

#### 4.1 | Conclusions, limitations, and future directions

This study presents data on traumatic experiences, ways of coping, and the potential for MBIs to help alleviate health disparities in an underserved, marginalised population. We note several limitations. Ours was a convenience sample of women from Southern California at a state-funded drug treatment facility, potentially limiting generalisability since treatment access and benefits vary by state and countries and men were not included. Focus groups were selected rather than interviews, although the literature on which is superior for discussing sensitive topics with vulnerable groups is not definitive (see Jordan et al., 2007 for a discussion). Data was collected immediately before the COVID-19 pandemic, thus due to statewide mitigation restrictions we were not able to return to site nor contact unhoused members of the CAB and focus groups to validate transcripts and data codes. All women had been in treatment for at least 30 days and do not represent perspectives from the acute phase of substance detox. Indeed, meta-analytic findings suggest mindfulness is not an appropriate first-line treatment for the acute phases of psychiatric conditions (Andersson et al., 2014). While our data provide a favourable view of the potential for mindfulness to benefit

trauma exposed women with co-occurring PTSD and SUD, such techniques may be most beneficial after obtaining some psychological stability or during the maintenance and relapse prevention phases of treatment. While trauma exposure was exceedingly high, women reported a variety of adaptive, approach based, and proactive coping techniques, as well as healthy coping skills related to emotion regulation. While there are important challenges to consider, on balance, mindfulness-based techniques may help promote abstinence and facilitate recovery in WEH. Taken together, our data provide an optimistic view of the potential for resilience and recovery in WEH who have experienced severe trauma and high health disparities.

#### ACKNOWLEDGEMENTS

We thank James Klotzle, Terri Hagmann, Maina Minchin, and the entire staff at Prototypes and Serenity House for their invaluable assistance with the data collection. We thank Kartik Yadav, MSCR for his expertise and assistance throughout the project. Finally, we thank the women in our study who shared their experience and strength with us. Work was supported by National Institute of Minority Health and Health Disparities MD013910 to Dana Rose Garfin. Dr. Garfin was at the University of California, Irvine at the time of data collection.

#### CONFLICT OF INTEREST

The authors have no conflict of interests to report.

#### DATA AVAILABILITY STATEMENT

To protect the privacy and confidentiality of the participants, redacted data is available upon request from the authors.

#### REFERENCES

- Alhusen, J. L., Norris-Shortle, C., Cosgrove, K., & Marks, L. (2017). I'm opening my arms rather than pushing away: Perceived benefits of a mindfulness-based intervention among homeless women and young children. *Infant Mental Health Journal*, 38(3), 434–442. <https://doi.org/10.1002/imhj.21641>
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). American Psychiatric Publishing.
- Andersson, G., Cuijpers, P., Carlbring, P., Riper, H., & Hedman, E. (2014). Internet-based vs. face-to-face cognitive behaviour therapy for psychiatric and somatic disorders: A systematic review and meta-analysis. *World Psychiatry*, 13(3), 1–23. <https://doi.org/10.1002/wps.20151>
- Ayano, G., Solomon, M., Tsegay, L., Yohannes, K., & Abraha, M. (2020). A systematic review and meta-analysis of the prevalence of post-traumatic stress disorder among homeless people. *Psychiatric Quarterly*, 91(4), 949–963. <https://doi.org/10.1007/s11126-020-09746-1>
- Ayer, L. A., Harder, V. S., Rose, G. L., & Helzer, J. E. (2011). Drinking and stress: An examination of sex and stressor differences using IVR-based daily data. *Drug and Alcohol Dependence*, 115(3), 205–212. <https://doi.org/10.1016/j.drugalcdep.2010.10.022>
- Baer, R. A. (2003). Mindfulness training as a clinical intervention: A conceptual and empirical review. *Clinical Psychology: Science and Practice*, 10(2), 125–143. <https://doi.org/10.1093/clipsy/bpg015>
- Bailey, K., Trevillion, K., & Gilchrist, G. (2019). What works for whom and why: A narrative systematic review of interventions for reducing post-traumatic stress disorder and problematic substance use among women with experiences of interpersonal violence. *Journal of*

- Substance Abuse Treatment*, 99, 88–103. <https://doi.org/10.1016/j.jsat.2018.12.007>
- Banyard, V. L., & Graham-Bermann, S. A. (1998). Surviving poverty: Stress and coping in the lives of housed and homeless mothers. *American Journal of Orthopsychiatry*, 68(3), 479–489. <https://doi.org/10.2307/j.ctt1bj4rs0>
- Bowen, S., De Bower, D., & Bergman, A. L. (2017). The role of mindfulness as approach-based coping in the PTSD-substance abuse cycle. *Addictive Behaviors*, 64, 212–216. <https://doi.org/10.1016/j.addbeh.2016.08.043>
- Bowen, S., Witkiewitz, K., Clifasefi, S. L., Grow, J., Chawla, N., Hsu, S. H., Carroll, H. A., Harrop, E., Collins, S. E., Lustyk, M. K., & Larimer, M. E. (2014). Relative efficacy of mindfulness-based relapse prevention, standard relapse prevention, and treatment as usual for substance use disorders. *JAMA Psychiatry*, 71(5), 547–556. <https://doi.org/10.1001/jamapsychiatry.2013.4546>
- Bower, J. E., Irwin, M. R., & Sciences, B. (2017). Mind-body therapies and control of inflammatory biology: A descriptive review. *Brain, Behavior, and Immunity*, 51, 1–11. <https://doi.org/10.1016/j.bbi.2015.06.012>
- Boxer, P., & Sloan-Power, E. (2013). Coping with violence: A comprehensive framework and implications for understanding resilience. *Trauma, Violence, & Abuse*, 14(3), 209–221. <https://doi.org/10.1177/1524838013487806>
- Boyd, J. E., Lanius, R. A., & McKinnon, M. C. (2018). Mindfulness-based treatments for posttraumatic stress disorder: A review of the treatment literature and neurobiological evidence. *Journal of Psychiatry & Neuroscience*, 43(1), 7–25. <https://doi.org/10.1503/jpn.170021>
- Breslin, F. C., Zack, M., & McMains, S. (2002). An information-processing analysis of mindfulness: Implications for relapse prevention in the treatment of substance abuse. *Clinical Psychology: Science and Practice*, 9(3), 275–299. <https://doi.org/10.1093/clipsy.9.3.275>
- California Department of Health Care Services. (2022). <https://www.dhcs.ca.gov/services/adp>
- Campbell, J. L., Quincy, C., Osserman, J., & Pedersen, O. K. (2013). Coding in-depth semistructured interviews: Problems of unitization and intercoder reliability and agreement. *Sociological Methods & Research*, 42(3), 294–320. <https://doi.org/10.1177/0049124113500475>
- Coenen, M., Stamm, T. A., Stucki, G., & Cieza, A. (2012). Individual interviews and focus groups in patients with rheumatoid arthritis: A comparison of two qualitative methods. *Quality of Life Research*, 21(2), 359–370. <https://doi.org/10.1007/s11136-011-9943-2>
- Davidson, L., Andres-Hyman, R., Bedregal, L., Tondora, J., Frey, J., & Kirk, T. A. (2008). From 'double trouble' to 'dual recovery': Integrating models of recovery in addiction and mental health. *Journal of Dual Diagnosis*, 4(3), 273–290. <https://doi.org/10.1080/15504260802072396>
- Duncan, S., Oby, S., & Larkin, H. (2019). Trauma and adversity in the lives of people experiencing homelessness. In *Homelessness prevention and intervention in social work* (pp. 41–56).
- Dutton, M. A., Bermudez, D., Matas, A., Majid, H., & Myers, N. L. (2014). Mindfulness-based stress reduction for low-income predominately african-american women with PTSD and a history of intimate partner violence. *Cognitive and Behavioral Practice*, 20(1), 23–32. <https://doi.org/10.1016/j.cbpra.2011.08.003>
- Fonfield-Ayinla, G. (2009). Commentary: A consumer perspective on parenting while homeless. *American Journal of Orthopsychiatry*, 79(3), 299–300. <https://doi.org/10.1037/a0017239>
- Glaser, B. G., & Strauss, A. L. (1967). *The discovery of grounded theory*. Aldine Publishing Company.
- Gray, M. J., Litz, B. T., Hsu, J. L., & Lombardo, T. W. (2004). Psychometric properties of the life events checklist. *Assessment*, 11(4), 330–341. <https://doi.org/10.1177/1073191104269954>
- Guest, G., Namey, E., Taylor, J., Eley, N., & McKenna, K. (2017). Comparing focus groups and individual interviews: Findings from a randomized study. *International Journal of Social Research Methodology*, 20(6), 693–708. <https://doi.org/10.1080/13645579.2017.1281601>
- Guillén, A. I., Marín, C., Panadero, S., & Vázquez, J. J. (2020). Substance use, stressful life events and mental health: A longitudinal study among homeless women in Madrid (Spain). *Addictive Behaviors*, 103, 106246. <https://doi.org/10.1016/j.addbeh.2019.106246>
- Gültekin, L., Brush, B. L., Baiardi, J. M., Kirk, K., & VanMaldeghem, K. (2014). Voices from the street: Exploring the realities of family homelessness. *Journal of Family Nursing*, 20(4), 390–414. <https://doi.org/10.1177/1074840714548943>
- Hayes, S. C., Strosahl, K., Wilson, K. G., Bissett, R. T., Pistorello, J., Toarmino, D., Polusny, M. A., Dykstra, T. A., Batten, S. V., Bergan, J., Stewart, S. H., Zvolensky, M. J., Eifert, G. H., Bond, F. W., Forsyth, J. P., Karekla, M., & Mccurry, S. M. (2004). Measuring experiential avoidance: A preliminary test of a working model. *The Psychological Record*, 54(4), 553–578. <https://doi.org/10.1007/BF03395492>
- Himelstein, S. (2011). Meditation research: The state of the art in correctional settings. *International Journal of Offender Therapy and Comparative Criminology*, 55(4), 646–661. <https://doi.org/10.1177/0306624X10364485>
- Hopper, E. K., Bassuk, E. L., & Olivet, J. (2010). Shelter from the storm: Trauma-informed care in homelessness services settings. *The Open Health Services and Policy Journal*, 3(2), 80–100.
- Israel, B. A., Schulz, A. J., Parker, E. A., & Becker, A. B. (2001). Community-based participatory research: Policy recommendations for promoting a partnership approach in health research. *Education and Health*, 14(2), 182–197.
- Ivankova, N., & Wingo, N. (2018). Applying mixed methods in action research: Methodological potentials and advantages. *American Behavioral Scientist*, 62(7), 978–997. <https://doi.org/10.1177/0002764218772673>
- Jenzer, T., Meisel, S. N., Blayney, J. A., Colder, C. R., & Read, J. P. (2020). Reciprocal processes in trauma and coping: Bidirectional effects over a four-year period. *Psychological Trauma: Theory, Research, Practice, and Policy*, 12(2), 207–218. <https://doi.org/10.1037/tra0000500>
- Jordan, J., Lynch, U., Moutray, M., O'Hagan, M.-T., Orr, J., Peake, S., & Power, J. (2007). Using focus groups to research sensitive issues: Insights from group interviews on nursing in the Northern Ireland Troubles. *International Journal of Qualitative Methods*, 6(4), 1–19. <https://doi.org/10.1177/160940690700600401>
- Kabat-Zinn, J. (1990). *Full catastrophe living: Using the wisdom of your body and mind to face stress, pain, and illness*. Bantam Books.
- Karadzov, D., Yuan, Y., & Bond, L. (2020). Coping amidst an assemblage of disadvantage: A qualitative metasynthesis of first-person accounts of managing severe mental illness while homeless. *Journal of Psychiatric and Mental Health Nursing*, 27(1), 4–24. <https://doi.org/10.1111/jpm.12524>
- Keyes, K. M., Hatzenbuehler, M. L., & Hasin, D. S. (2011). Stressful life experiences, alcohol consumption, and alcohol use disorders: The epidemiologic evidence for four main types of stressors. *Psychopharmacology*, 218(1), 1–17. <https://doi.org/10.1007/s00213-011-2236-1>
- Korecki, J. R., Schwebel, F. J., Votaw, V. R., & Witkiewitz, K. (2020). Mindfulness-based programs for substance use disorders: A systematic review of manualized treatments. *Substance Abuse Treatment, Prevention, and Policy*, 15(1), 51. <https://doi.org/10.1186/s13011-020-00293-3>
- Lafavor, T., Boer, D. De, & Poole, M. (2020). Intergenerational effects of early parental adversity on child developmental outcomes among families living in emergency homeless shelters. *Journal of Health Care for the Poor and Underserved*, 31(3), 1264–1280. <https://doi.org/10.1353/hpu.2020.0093>

- Lang, A. J., Strauss, J. L., Bomyea, J., Bormann, J. E., Hickman, S. D., Good, R. C., & Essex, M. (2012). The theoretical and empirical basis for meditation as an intervention for PTSD. *Behavior Modification*, 36(6), 759–786. <https://doi.org/10.1177/0145445512441200>
- Lazarus, R. S., & Folkman, S. (1984). *Stress, appraisal, and coping*. Springer publishing company.
- Linneberg, M. S., & Korsgaard, S. (2019). Coding qualitative data: A synthesis guiding the novice. *Qualitative Research Journal*, 19(3), 259–270. <https://doi.org/10.1108/QRJ-12-2018-0012>
- Maddock, A., Hevey, D., & Eidenmueller, K. (2017). Mindfulness training as a clinical intervention with homeless adults: A pilot study. *International Journal of Mental Health and Addiction*, 15(3), 529–544. <https://doi.org/10.1007/s11469-016-9718-7>
- Manning, J. (2017). *In vivo coding*. *The International Encyclopedia of Communication Research Methods*, 1–2. <https://doi.org/10.1002/9781118901731.iecrm0270>
- Matthews, A. K., Anderson, E. E., Willis, M., Castillo, A., & Choure, W. (2018). A Community Engagement Advisory Board as a strategy to improve research engagement and build institutional capacity for community-engaged research. *Journal of Clinical and Translational Science*, 2(2), 66–72. <https://doi.org/10.1017/cts.2018.14>
- Maxwell, J. A., Chmiel, M., & Rogers, S. E. (2015). Designing integration in multimethod and mixed methods research. In S. N. Hesse-Biber & R. B. Johnson (Eds.), *The oxford handbook of multimethod and mixed methods research inquiry*. <https://doi.org/10.1093/oxfordhb/9780199933624.013.16>
- McMordie, L. (2021). Avoidance strategies: Stress, appraisal and coping in hostel accommodation. *Housing Studies*, 36(3), 380–396. <https://doi.org/10.1080/02673037.2020.1769036>
- Mills, K. L. (2009). 'Between pain and nothing, I choose nothing: Trauma, post-traumatic stress disorder and substance use. *Addiction*. <https://doi.org/10.1111/j.1360-0443.2009.02675.x>
- North, C. S., & Smith, E. M. (1992). Posttraumatic stress disorder among homeless men and women. *Psychiatric Services*, 43(10), 1010–1016. <https://doi.org/10.1176/ps.43.10.1010>
- Nyamathi, A. (1989). Comprehensive health seeking and coping paradigm. *Journal of Advanced Nursing*, 14(4), 281–290. <https://doi.org/10.1111/j.1365-2648.1989.tb03415.x>
- Nyamathi, A., Salem, B. E., Shin, S. S., Jones, A. A., Garfin, D. R., Yadav, K., Chang, A. H., White, K., & Morisky, D. (2021). Effect of a nurse-led community health worker intervention on latent tuberculosis medication completion among homeless adults. *Nursing Research*, 70(6), 433–442. <https://doi.org/10.1097/NNR.0000000000000545>
- Nyamathi, A. M., Stein, J. A., & Bayley, L. J. (2000). Predictors of mental distress and poor physical health among homeless women. *Psychology and Health*, 15(4), 483–500. <https://doi.org/10.1080/08870440008402008>
- Owen, S. (2001). The practical, methodological and ethical dilemmas of conducting focus groups with vulnerable clients. *Journal of Advanced Nursing*, 36(5), 652–658. <https://doi.org/10.1046/j.1365-2648.2001.02030.x>
- Padgett, D. K., Smith, B. T., Henwood, B. F., & Tiderington, E. (2012). Life course adversity in the lives of formerly homeless persons with serious mental illness: Context and meaning. *American Journal of Orthopsychiatry*, 82(3), 421–430. <https://doi.org/10.1111/j.1939-0025.2012.01159.x>
- Phipps, M., Dalton, L., Maxwell, H., & Cleary, M. (2019). Women and homelessness, a complex multidimensional issue: Findings from a scoping review. *Journal of Social Distress and the Homeless*, 28(1), 1–13. <https://doi.org/10.1080/10530789.2018.1534427>
- Polusny, M. A., Erbes, C. R., Thuras, P., Moran, A., Lamberty, G. J., Collins, R. C., Rodman, J. L., & Lim, K. O. (2015). Mindfulness-based stress reduction for posttraumatic stress disorder among veterans a randomized clinical trial. *JAMA*, 314(5), 456–465. <https://doi.org/10.1001/jama.2015.8361>
- Prins, A., Bovin, M. J., Kimerling, R., Kaloupek, D. G., Marx, B. P., Pless Kaiser, A., & Schnurr, P. P. (2015). Primary care PTSD screen. *Measurement Instrument*.
- Prins, A., Bovin, M. J., Smolenski, D. J., Mark, B. P., Kimerling, R., Jenkins-Guarnieri, M. A., Kaloupek, D. G., Schnurr, P. P., Pless Kaiser, A., Leyva, Y. E., & Tiet, Q. Q. (2016). The primary care PTSD screen for DSM-5 (PC-PTSD-5): Development and evaluation within a veteran primary care sample. *Journal of General Internal Medicine*, 31, 1206–1211. <https://doi.org/10.1007/s11606-016-3703-5>
- Roberts, N. P., Roberts, P. A., Jones, N., & Bisson, J. I. (2015). Psychological interventions for post-traumatic stress disorder and comorbid substance use disorder: A systematic review and meta-analysis. *Clinical Psychology Review*, 38, 25–38. <https://doi.org/10.1016/j.cpr.2015.02.007>
- Saxena, P., Messina, N. P., & Grella, C. E. (2014). Who benefits from gender-responsive treatment?: Accounting for abuse history on longitudinal outcomes for women in prison. *Criminal Justice and Behavior*, 41(4), 417–432. <https://doi.org/10.1177/0093854813514405>
- Schmitt, M., Dressel, A., Del Carmen Graf, M., Pittman, B., Deal, E., Krueger, E., Lopez, A. A., Kako, P., & Mkandawire-Valhmu, L. (2021). Adverse childhood experiences among previously homeless African American women. *Public Health Nursing*, 39(2), 1–10. <https://doi.org/10.1111/phn.12970>
- Schouten, K. A., van Hooren, S., Knipscheer, J. W., Kleber, R. J., & Hutchemaekers, G. J. M. (2019). Trauma-focused art therapy in the treatment of posttraumatic stress disorder: A pilot study. *Journal of Trauma & Dissociation*, 20(1), 114–130. <https://doi.org/10.1080/15299732.2018.1502712>
- Sheerin, C. M., Chowdhury, N., Lind, M. J., Kurtz, E. D., Rappaport, L. M., Berenz, E. C., Brown, R. C., Pickett, T., McDonald, S. D., Danielson, C. K., & Amstadter, A. B. (2018). Relation between coping and posttrauma cognitions on PTSD in a combat-trauma population. *Military Psychology*, 30(2), 98–107. <https://doi.org/10.1080/08995605.2017.1420980>
- Sibinga, E. M. S., Kerrigan, D., Stewart, M., Johnson, K., Magyari, T., & Ellen, J. M. (2011). Mindfulness-based stress reduction for urban youth. *Journal of Alternative & Complementary Medicine*, 17(3), 213–218. <https://doi.org/10.1089/acm.2009.0605>
- Suls, J., & Fletcher, B. (1985). The relative efficacy of avoidant and non-avoidant coping strategies: A meta-analysis. *Health Psychology*, 4(3), 249–288. <https://doi.org/10.1037/0278-6133.4.3.249>
- Thompson, N. J., Fiorillo, D., Rothbaum, B. O., Ressler, K. J., & Michopoulos, V. (2018). Coping strategies as mediators in relation to resilience and posttraumatic stress disorder. *Journal of Affective Disorders*, 225, 153–159. <https://doi.org/10.1016/j.jad.2017.08.049>
- Thompson, R. W., Arnkoff, D. B., & Glass, C. R. (2011). Conceptualizing mindfulness and acceptance as components of psychological resilience to trauma. *Trauma, Violence, & Abuse*, 12(4), 220–235. <https://doi.org/10.1177/1524838011416375>
- Tsai, A. C., Weiser, S. D., Dilworth, S. E., Shumway, M., & Riley, E. D. (2015). Violent victimization, mental health, and service utilization outcomes in a cohort of homeless and unstably housed women living with or at risk of becoming infected with HIV. *American Journal of Epidemiology*, 181(10), 817–826. <https://doi.org/10.1093/aje/kwu350>
- Tsai, J., Rosenheck, R. A., Decker, S. E., Desai, R. A., & Harpaz-Rotem, I. (2012). Trauma experience among homeless female veterans: Correlates and impact on housing, clinical, and psychosocial outcomes. *Journal of Traumatic Stress*, 25(6), 624–632. <https://doi.org/10.1002/jts.21750>
- Upshur, C. C., Jenkins, D., Weinreb, L., Gelberg, L., & Orvek, E. A. (2017). Prevalence and predictors of substance use disorders among homeless women seeking primary care: An 11 site survey. *American Journal on Addictions*, 26(7), 680–688. <https://doi.org/10.1111/ajad.12582>

- van Zuiden, M., Kavelaars, A., Geuze, E., Olf, M., & Heijnen, C. J. (2013). Predicting PTSD: Pre-existing vulnerabilities in glucocorticoid-signaling and implications for preventive interventions. *Brain, Behavior, and Immunity*, 30, 12–21. <https://doi.org/10.1016/j.bbi.2012.08.015>
- Whitzman, C. (2006). At the intersection of invisibilities: Canadian women, homelessness and health outside the “big city. *Gender, Place & Culture*, 13(4), 383–399. <https://doi.org/10.1080/09663690600808502>
- Williams, M., & Moser, T. (2019). The art of coding and thematic exploration in qualitative research. *International Management Review*, 15(1), 45–55.
- Witkiewitz, K., Marlatt, G. A., & Walker, D. (2005). Mindfulness-Based Relapse Prevention for alcohol and substance use disorders. *Journal of Cognitive Psychotherapy: International Quarterly*, 19(3), 211–228. <https://doi.org/10.1891/jcop.2005.19.3.211>
- Wu, G., Feder, A., Cohen, H., Kim, J. J., Calderon, S., Charney, D. S., & Mathé, A. A. (2013). Understanding resilience. *Frontiers in Behavioral Neuroscience*, 7, 1–15. <https://doi.org/10.3389/fnbeh.2013.00010>
- Yehuda, R., Daskalakis, N. P., Bierer, L. M., Bader, H. N., Klengel, T., Holsboer, F., & Binder, E. B. (2016). Holocaust exposure induced

intergenerational effects on FKBP5 methylation. *Biological Psychiatry*, 80(5), 372–380. <https://doi.org/10.1016/j.biopsych.2015.08.005>

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Garfin, D. R., Amador, A., Osorio, J., Ruivivar, K. S., Torres, A., & Nyamathi, A. M. (2022). A multi-method exploration of mindfulness as a coping tool: Perspectives from trauma-exposed, unhoused women residing at a drug treatment facility. *Stress and Health*, 1–14. <https://doi.org/10.1002/smi.3188>