

Acute Stress, Worry, and Impairment in Health Care and Non-Health Care Essential Workers During the COVID-19 Pandemic

Dana Rose Garfin^{1, 2}, Lindita Djokovic³, Roxane Cohen Silver^{2, 4, 5}, and E. Alison Holman^{1, 4}

¹ Sue & Bill Gross School of Nursing, University of California, Irvine

² Program in Public Health, University of California, Irvine

³ Department of Counseling, Clinical, and School Psychology, University of California, Santa Barbara

⁴ Department of Psychological Science, University of California, Irvine

⁵ Department of Medicine, University of California, Irvine


Objective: Health care and non-health care essential workers working in face-to-face interactions during the coronavirus disease 2019 (COVID-19) pandemic may be vulnerable to psychosocial distress. Limited empirical research on COVID-19-related psychosocial outcomes has utilized probability-based samples including both health care and non-health care essential workers. **Method:** We surveyed a sample of 1,821 United States self-identified essential workers, collected using probability-based methods, working in face-to-face interactions during the early phase of the COVID-19 outbreak (March 18, 2020 through April 18, 2020), in three consecutive 10-day cohorts. We assessed acute stress, health-related worries, and functional impairment. Demographics, secondary stressors (lack of childcare or health care, lost wages), and pre-COVID-19 mental and physical health were examined as predictors of psychological outcomes. **Results:** Acute stress ($\beta = .08, p = .001$), health-related worries ($\beta = .09, p = .001$), and functional impairment ($\beta = .05, p = .034$) increased over time in the early weeks of the outbreak. Health care essential workers reported lower functional impairment ($\beta = -.06, p = .009$) and acute stress ($\beta = -.06, p = .015$) compared with non-health care essential workers. Across the sample, prior mental and physical health ailments, inability to obtain health care, lost wages, younger age, female gender, and Hispanic ethnicity were associated with acute stress ($\beta s = -.14$ to $.15, ps \leq .001$), health-related worries ($\beta s = -.09$ to $.14, ps \leq .001$), and functional impairment ($\beta s = -.08$ to $.16, ps \leq .006$). Lack of childcare ($\beta = .09, p < .001$) was positively associated with acute stress. **Conclusion:** Non-health care essential workers may be vulnerable to negative psychosocial outcomes. Targeted training and support may help facilitate coping with the effects of working in-person during the ongoing COVID-19 pandemic. Findings may help inform intervention efforts, critical as COVID-19 becomes endemic and society must learn to live with its evolving variants.


Clinical Impact Statement


Prior literature has documented health care essential workers' adverse psychological responses during viral outbreaks including coronavirus disease 2019 (COVID-19); non-health care essential workers have been understudied. Between March 18, 2020 and April 18, 2020, we surveyed 1,821 United States residents (a subsample of a nationally representative sample of 6,514) still working in-person during the COVID-19 pandemic. Demographics, secondary stressors, non-health care occupation, and mental/physical health history were positively associated with acute stress, health-related worries, and functional impairment. As society adapts to COVID-19 and its evolving variants, essential workers may continue to be vulnerable. Adequate protection, targeted psychosocial services, and appropriate preparation for future outbreaks is crucial.

Keywords: COVID-19, acute stress, essential workers, mental health, functional impairment

Supplemental materials: <https://doi.org/10.1037/tra0001224.supp>

Dana Rose Garfin  <https://orcid.org/0000-0002-0435-9307>

Roxane Cohen Silver  <https://orcid.org/0000-0002-4013-6792>

E. Alison Holman  <https://orcid.org/0000-0001-5076-8403>

Research was funded by National Science Foundation RAPID Grant SES 2026337. Dana Rose Garfin was supported by K01 MD013910. The

study sponsor played no role in study design; collection, analysis, and interpretation of data; writing the report; and the decision to submit the report for publication. Authors report no conflicts of interest.

Correspondence concerning this article should be addressed to Dana Rose Garfin, 100C Berk Hall, Irvine, CA 92697, United States. Email: dgarfin@uci.edu

On March 13, 2020, the White House declared the coronavirus disease 2019 (COVID-19) outbreak a national emergency, its protracted duration and associated psychosocial impact unforeseeable. State officials issued sweeping stay-at-home orders that exempted essential workers, including health care workers, caregivers, and grocery store employees, as well as workers in other sectors whose jobs required in-person interactions with social distancing guidelines difficult or impossible to maintain (Lai et al., 2020; Shanafelt et al., 2020). The empirical literature on psychosocial responses to COVID-19 among essential workers has primarily focused on caregivers and health care workers (Lai et al., 2020; Shanafelt et al., 2020), largely neglecting the experiences of non-health care essential workers.

Even during the most stringent COVID-19-related restrictions, many essential workers continued in-person work, often due to lack of job security or paid leave, and sometimes without health insurance, amplifying COVID-19-relevant worries about one's health and safety (Shanafelt et al., 2020). Despite the increasing availability of vaccines, COVID-19 has remained a threat, due to emerging variants such as the Delta and Omicron variants (Christie et al., 2021) and vaccine hesitancy (Kofman et al., 2021). As COVID-19 becomes endemic, those on the front lines will be forced to adapt and respond to the ongoing threat of COVID-19 (Kofman et al., 2021).

Research conducted during COVID-19 has demonstrated that health care workers are at risk for psychosocial maladies (Benfante et al., 2020), particularly those working on the front lines (Lai et al., 2020). While this relationship has been demonstrated during prior viral outbreaks (Cabarkapa et al., 2020), during COVID-19 these concerns have been exacerbated for non-health care, public-facing occupations as well: like health care workers, essential workers from other industries are at increased risk for COVID-19 infection due to frequent interactions with the public (Baker et al., 2020). Thus, there has been a growing concern to manage the physical and psychological health of all frontline workers (Sim, 2020), critical for effective workplace performance (Wright & Cropanzano, 2000) when essential tasks must be sustained. In summary, it is important to understand the psychosocial experience of workers from both health care and non-health care sectors critical for societal functioning during the COVID-19 pandemic and as society prepares for future viral threats, which science suggest will increase in the years ahead (Rogalski et al., 2017).

Health Care Essential Workers

Research on psychosocial outcomes in health care essential workers during COVID-19 has proliferated, yet a clearer understanding of specific risk factors associated with psychosocial maladies may further guide research and recommendations. Meta-analytic findings regarding mental health in health care workers (including doctors, nurses, and other medical personnel) during the early phase of COVID-19 (e.g., before mid-April, 2020) found a 23% prevalence of clinically significant anxiety, although these rates were comparable with that of the general public, and at the lower end of what was reported in health care workers during the Middle East respiratory syndrome (MERS) and severe acute respiratory syndrome (SARS) outbreaks (Pappa et al., 2020). In a rapid review of COVID-19's impact on health care worker's mental health, concerns over family health and safety, in addition to fear of infection, were risk factors for adverse mental health outcomes

(De Kock et al., 2021). A recent literature review of the mental health of frontline health care workers employed during viral outbreaks revealed that persistent, generalized stress was associated with subsequent downstream problems including panic attacks, insomnia, and burnout (Magill et al., 2020). Internationally, health care essential workers who continued working during the pandemic also screened positive for moderate to extremely severe stress, psychological distress (Chew, Lee, et al., 2020), and post-traumatic stress disorder (PTSD; Tan et al., 2021). While data has illustrated that factors associated with patient care (e.g., direct work with COVID-19 patients) are associated with psychological burdens such as anxiety in health care workers (Lai et al., 2020), few samples have been drawn from representative samples or included comparisons with non-health care essential workers.

Research on health care worker mental health during prior viral epidemics revealed mental health ailments like acute stress disorder increased in frequency, particularly when certain sociodemographic (i.e., younger age and female gender) and occupational (i.e., high-risk roles) factors were present (Serrano-Ripoll et al., 2020). More recently, history of poor physical health (Chew, Ngiam, et al., 2020; De Kock et al., 2021; Tan et al., 2020) and prior psychiatric diagnoses (Elbay et al., 2020) have been recognized as additional risk factors for COVID-19-related psychological distress in health care workers, which may have implications for present functioning and downstream physical and mental health (Garfin et al., 2018).

Non-Health Care Essential Workers

During strict stay-at-home restrictions in the United States during Spring, 2020, many non-health care essential workers (e.g., food service or transportation workers) continued to engage with the public, at times with little work flexibility and high risk of contracting COVID-19 (Sim, 2020). Such employees often work with less stringent protective protocols (Steege et al., 2009) than health care sector workers. As the economy remains open, vaccine hesitancy persists (Sallam, 2021), vaccination rates are well-below targets, and new variants impose increasing risk to frontline works (Christie et al., 2021), understanding predictors of distress and impairment in essential workers is critical. Global research using non-probability-based samples found non-health care workers have continually reported adverse psychosocial outcomes during the COVID-19 pandemic. For example, up to 50% of Indian migrant workers screened positive for anxiety (Grover et al., 2021) and 65.1% of Spanish non-health care essential workers endorsed psychological distress (i.e., constantly feeling overwhelmed and stressed) during the early phase of the COVID-19 pandemic (Ruiz-Frutos et al., 2021). Moreover, non-health care essential workers have also reported health-related worries, with loved ones dying from COVID-19 and the health and well-being of family or loved ones indicated as top COVID-19-related concerns (Toh et al., 2021). Research using probability-based samples could strengthen inferences and elucidate key predictors of such maladies.

Some empirical literature suggests that non-health care essential workers actively employed during the COVID-19 pandemic are also at risk for psychological maladies, and in some cases potentially at greater risk than health care workers. Recent research on COVID-19 has demonstrated that some of those in non-health care occupations (i.e., agricultural workers) report higher generalized

distress symptom scores compared with other groups, including those working in health care industries (Tian et al., 2020). Similarly, in a sample of Australian workers, non-health care workers reported significantly higher levels of anxiety, stress, and significantly lower quality of life compared with health care workers (Toh et al., 2021). Yet, a community-based sample in Turkey indicated health care workers were more distressed than the general public (Hacimusalar et al., 2020). The present study seeks to extend this work in a U.S. sample by using a methodologically rigorous design (i.e., probability-based sampling), drawing from prior literature on disaster psychology to account for additional risk factors.

Secondary Stressors

Secondary stressors are individual-level events (e.g., personal injury or illness, loss of a job) that occur as part of a disaster cascade (Garfin et al., 2014). Exposure to greater numbers of secondary stressors has been associated with greater psychological distress following prior collective traumas including earthquakes (Garfin et al., 2014), hurricanes (Galea et al., 2007), and other natural disasters (Kessler et al., 2012). Less is known about how exposure to secondary stressors during a collective trauma like the COVID-19 pandemic may exacerbate psychosocial responses, particularly among essential workers. Data collected during the early phase of COVID-19 found some secondary stressors occurred for a substantial minority of respondents; for example, inability to obtain health care was associated with serious psychological distress (i.e., feeling nervous, restless, or hopeless; McGinty et al., 2020). In a community-based sample that included health care and non-health care workers, secondary stressors including difficulties finding childcare were positively associated with anxiety (Hacimusalar et al., 2020). Early research also highlighted economic stressors as an additional risk factor for distress, such as financial loss, which during the early phases of the pandemic appeared to amplify psychological distress in the general population (Zheng et al., 2021). As the COVID-19 pandemic persisted, those unable to work from home may have been vulnerable to these stressors, which may have correlated with adverse psychosocial outcomes.

Preexisting Mental and Physical Health Vulnerabilities

Elevated risk of severe COVID-19-related complications and mortality for individuals with preexisting physical health conditions has been highly publicized (Flaherty et al., 2020), potentially eliciting worry, distress, and impairment in those with such vulnerabilities. This was evident during prior viral outbreaks: in a systematic review on the impact of viral epidemics on mental health outcomes, chronic illness was identified as a high-risk factor for experiencing psychiatric symptoms (Luo et al., 2020). Pre-event mental health diagnoses (e.g., anxiety disorders and mood disorders) may also portended greater problems: for example, prior mental health ailments were associated with psychological distress and functional impairment following the 2014 Ebola public health crisis (Thompson et al., 2017). During the COVID-19 pandemic specifically, such ailments were associated with COVID-19 related stress and traumatic stress symptoms among American and Canadian adults (Asmundson et al., 2020). In health care workers, pre-COVID-19 psychological treatment increased the risk for developing COVID-19-related posttraumatic distress and anxiety by 2.5

times (Asmundson et al., 2020). These findings demonstrate the importance of evaluating preexisting physical and mental health ailments as potential risk factors for maladaptive psychosocial outcomes during COVID-19.

Demographic Risk Factors

Demographic risk factors may also correlate with psychological distress and impairment in essential workers, potentially exacerbating existing health disparities (Kantamneni, 2020). For example, racial/ethnic minorities often encounter more COVID-19 related occupational inequalities (e.g., high-exposure occupations) and discrimination (Kantamneni, 2020), potentially leading to worse mental health outcomes. Lower income negatively correlates with deleterious psychosocial outcomes; for example, individuals who lost income due to COVID-19 reported higher hopelessness and anxiety compared with those who did not lose income (Hacimusalar et al., 2020). Low educational attainment may also place low-wage workers at more physical risk during COVID-19 (Gallagher et al., 2021). Finally, research demonstrates female health care essential workers face higher risks for maladaptive mental health outcomes during COVID-19 compared to their male counterparts (Cabarkapa et al., 2020; Elbay et al., 2020; Lai et al., 2020; Pappa et al., 2020).

Other demographic indicators may also portend increased vulnerability to psychosocial maladies. Despite reports that older frontline workers actively working during COVID-19 were more vulnerable to death and infection (Ghilarducci & Farmand, 2020), research has consistently shown younger workers report higher psychological distress compared to older workers (Ruiz-Frutos et al., 2021). Regional variation, a potential proxy for level of community transmission and severity of the outbreak (e.g., hospitalization and death rates), may also differentially predict psychosocial distress. Specifically, as the epidemic first spread across the United States, medical personnel in the New York Metropolitan Area were overwhelmed due to an influx of infected individuals and a lack of resources to support them (Konda et al., 2020).

The Present Study

As society continues to struggle to protect workers during the ongoing and continually evolving COVID-19 pandemic (Kofman et al., 2021), it is critical to understand experiences of essential workers who engage with the public and are at increased risk for infection. As such, the objective of the present study was to examine key theoretically and empirically derived predictors of psychosocial responses during COVID-19 among a sample of U.S. health care and non-health care essential workers still working in face-to-face interaction in the first months of the pandemic. We sought to improve upon the limited extant research by utilizing a probability-based sample, collected as the pandemic escalated in the United States. We had one hypothesis and one exploratory aim. We hypothesized that preexisting physical and mental health ailments and key secondary stressors would be associated with greater health-related worry, functional impairment, and acute stress, controlling for demographic risk factors. Given prior mixed research, as an exploratory aim we examined whether health care essential workers would report worse outcomes than non-health care essential workers.

Method

Sample and Procedure

Our study was conducted among a national sample of Americans using the NORC AmeriSpeak Panel, a probability-based panel of 35,000 U.S. households. NORC's AmeriSpeak Panel randomly selected participants from their panel to form a representative sample of U.S. households. Sample stratification promoted representativeness for age, gender, race/ethnicity, and education. NORC notified participants via e-mail when the confidential, 20-min survey was available. Three consecutive cohorts of panelists participated in the study. There was a 10-day fielding period among each of three cohorts: Cohort 1 (March 18, 2020 through March 28, 2020), Cohort 2 (March 29, 2020 through April 7, 2020), and Cohort 3 (April 8, 2020 through April 18, 2020). In total, 6,514 participants completed the survey (58.5% completion rate) across all cohorts with 85% responding within 3 days through computers (44%), smartphones (54%), and tablets (2%). Some procedures helped ensure valid survey responses. Respondents were removed from the main survey if they completed the survey in under one-third of the median duration and/or skipped more than 50% of the questions shown to them. Three respondents from Cohort 2 were removed since they completed the survey during the Cohort 1 timeframe and were not counted toward the total number of interviews delivered. All procedures were approved by the Institutional Review Board of University of California, Irvine; informed consent was obtained. A small compensation was provided (cash equivalent \$4). See online supplemental data for full list of measures used in these analyses.

From the full sample, a subsample of health care and non-health care essential workers was identified based on responses to several items. First, a subsample of health care workers was identified. As part of the profile data collected by NORC, panelist identify their employment type based on U.S. Bureau of Labor statistics identified 24 categories, including "Healthcare Practitioners and Technical Occupations." Occupations include physicians, paramedics, psychiatrists, nurses, surgical assistants, and physician assistants. (Full list of included occupations is available at <https://www.bls.gov/oes/current/oes290000.htm>.) Next, participants reported whether their job required in-person interaction and they were still working ($n = 1,525$). Participants also reported whether they were an essential worker asked to work extra hours ($n = 981$). Six hundred eighty-five participants endorsed both items, generating a final sample of "essential workers" ($n = 1,821$), of which 9.28% ($n = 169$) identified as health care workers.

Independent Variables

Secondary Stressors

Three items assessed COVID-19 related secondary stressors: lost wages, needing childcare due to COVID-19-related school closures, and inability to obtain health care due to COVID-19. Items were coded 0 (*not experienced*) or 1 (*experienced*).

Prior Physical and Mental Health Ailments

NORC collected health information history from participants before January 2020, before the official U.S. COVID-19 outbreak

announcement. Respondents were asked whether they had been diagnosed with physical health ailments (i.e., diagnoses of high cholesterol, hypertension, diabetes/high blood sugar, heart disease, stroke, cancer, lung disease, and other diagnoses). Prior physical health diagnoses were summed into an eight-item count variable. Respondents also reported previous mental health diagnoses (anxiety, depression, or any other emotional, nervous, or psychiatric diagnosis). Prior mental health diagnoses were coded 0 (*none*) or 1 (*prior anxiety, depression, or other emotional/nervous/psychiatric diagnosis*). These variables have been used in prior research (Holman et al., 2020).

Demographics

NORC collected demographic information (age, race/ethnicity, education, gender, income, and geographic region of residence) when participants enrolled into in the AmeriSpeak panel (updated annually for accuracy). Race/ethnicity was coded White, Black (non-Hispanic), other (two or more, non-Hispanic), and Hispanic. Education was coded less than high school, high school diploma, some college, and bachelor's degree and up. Income was classified as one of eight categories, consistent with prior analyses (Holman et al., 2020). Geographic regions of residence included Northeast, Midwest, South, and West.

Dependent Variables

Health-Related Worries

Health-related worries were assessed by asking participants how often they "Had fears that you will get very sick or die from the Coronavirus outbreak?" and "Worried that someone you care about (e.g., family, close friends) will get very sick or die from the Coronavirus outbreak?" Items were assessed using a 5-point scale from 1 (*never*) to 5 (*all the time*). Internal consistency was very good ($\alpha = .81$).

Functional Impairment

Prior week functional impairment was assessed using four items modified from the SF-36 that measured physical and emotional impairment (Ware & Sherbourne, 1992), on a 5-point scale from 1 (*none of the time*) to 5 (*all of the time*). Internal consistency was very good ($\alpha = .87$).

COVID-19-Related Acute Stress

COVID-19 outbreak-related acute stress symptoms were assessed using a 10-item modified version of the Acute Stress Disorder Scale 5 (Bryant, 2016). Items assessed symptoms from the previous week via a 5-point scale from 1 (*none at all*) to 5 (*a great deal*). Internal consistency was very good ($\alpha = .87$).

Statistical Analysis

All analyses were conducted using STATA 16. Bivariate comparisons (t tests for continuous variables and chi-square tests for categorical variables) illustrated significant differences between health care and non-health care essential workers. Three multivariate Ordinary Least Squares (OLS) regression analyses examined predictors of COVID-19-related health-related worries, functional impairment, and acute stress. Predictors were: health care worker status (0 = *non-health care essential worker*; 1 = *health*

care essential worker), secondary stressors (lack of childcare due to COVID-19 school closures, lack of health care, lost wages; each coded 0 = *did not occur*; 1 = *occurred*), demographics (i.e., age, ethnicity, education, female gender, and income), residential region, and cohort. Because the amount of missing data was extremely low (less than 1.5% in covariate-adjusted models) and Little’s Missing Completely at Random test was not significant ($\chi^2[9] = 4.19, p = .89$), listwise deletion was implemented. Post hoc, interaction terms between key predictors and essential worker status (health care or non-health care) were examined; each was examined separately in

individual models to conserve sample size, although results were identical when added into a combined model with all interaction terms.

Results

Table 1 presents descriptive statistics and between-group differences for demographic indicators and key covariates. Among all essential workers, difficulty meeting childcare needs was reported by 9.37% ($n = 169$); 9.26% ($n = 167$) reported inability

Table 1
Characteristics of the Sample (N = 1,821)

Essential worker characteristics	Health care essential workers (n = 169)	Non-health care essential workers (n = 1,652)	Test of between-group differences (non-health care vs. health care essential workers)	Effect size Cohen’s d (95% CI) or Cramer’s V
	<i>M (SD)</i>	<i>M (SD)</i>		
Age	42.37 (12.96)	43.88 (13.27)	$t(1,819) = 1.41$	$d = 0.11 [-0.05, 0.27]$
	<i>n (%)</i>	<i>n (%)</i>		
Race/ethnicity				
White, non-Hispanic	128 (75.74%)	1,181 (71.49%)		
Black, non-Hispanic	16 (9.47%)	170 (10.29%)		
Other/2+ races, non-Hispanic	12 (7.10%)	139 (8.41%)		
Hispanic	13 (7.69%)	162 (9.81%)	$\chi^2(3) = 1.52$	$V = 0.03$
Education				
Less than high school	0 (0%)	37 (2.24%)		
High school graduate	5 (2.96%)	279 (16.89%)		
Some college	58 (34.32%)	750 (45.40%)		
BA and above	106 (62.72%)	586 (35.47%)	$\chi^2(3) = 57.06^{**}$	$V = 0.18$
Gender				
Male	44 (26.04%)	751 (45.46%)		
Female	125 (73.96%)	901 (54.54%)	$\chi^2(1) = 23.52^{**}$	$V = 0.11$
Income				
Less than \$10,000	2 (1.18%)	66 (4.00%)		
\$10,000–\$24,999	5 (2.96%)	191 (11.56%)		
\$25,000–\$49,000	21 (12.43%)	403 (24.39%)		
\$50,000–\$74,999	40 (23.67%)	356 (21.55%)		
\$75,000–\$99,999	32 (18.93%)	272 (16.46%)		
\$100,000–\$124,999	39 (23.08%)	207 (12.53%)		
\$125,000–\$149,999	15 (8.88%)	89 (5.39%)		
\$150,000+	15 (8.88%)	68 (4.12%)	$\chi^2(7) = 47.61^{**}$	$V = 0.16$
Region				
Northeast	37 (21.89%)	224 (13.56%)		
Midwest	48 (28.40%)	466 (28.21%)		
South	45 (26.63%)	550 (33.29%)		
West	39 (23.08%)	412 (24.94%)	$\chi^2(3) = 9.73^*$	$V = 0.07$
Cohort				
Cohort 1: March 18, 2020 through March 28, 2020	57 (33.73%)	587 (35.53%)		
Cohort 2: March 29, 2020 through April 7, 2020	61 (36.09%)	533 (32.26%)		
Cohort 3: April 8, 2020 through April 18, 2020	51 (30.18%)	532 (32.20%)	$\chi^2(2) = 1.03$	$V = 0.02$
Secondary stressors				
Needed childcare for child because they were out of school	19 (11.38%)	150 (9.17%)	$\chi^2(1) = 0.87$	$V = 0.02$
I lost wages	51 (30.54%)	345 (21.09%)	$\chi^2(1) = 7.90^*$	$V = 0.07$
Unable to obtain necessary health care	12 (7.19%)	155 (9.47%)	$\chi^2(1) = 0.94$	$V = -0.02$
Prepandemic mental health problems	23 (13.61%)	282 (17.07%)	$\chi^2(1) = 1.31$	$V = -0.03$
	<i>M (SD)</i>	<i>M (SD)</i>		
Prepandemic physical health problems	0.65 (0.89)	0.83 (1.03)	$t(1,819) = 2.15^*$	$d = 0.17 [0.01, 0.33]$

Note. CI = confidence interval. Cohen’s *d* is presented for *t* tests; Cramer’s *V* for χ^2 . Prepandemic mental health problems = emotional/nervous/psychiatric problem, anxiety disorder, and depression (0 = none; 1 = one or more). Prepandemic physical health problems = high cholesterol, hypertension, diabetes/high blood sugar, heart disease, stroke, cancer, lung disease, and other diagnoses (0–8).

* $p < .05$. ** $p < .001$.

This document is copyrighted by the American Psychological Association or one of its allied publishers. This article is intended solely for the personal use of the individual user and is not to be disseminated broadly.

to obtain necessary health care; and 21.96% ($n = 396$) reported COVID-19-related lost or reduced wages. The means for acute stress, health-related worries, and functional impairment were 1.87 ($SD = .72$, range = 1–4.9), 2.53 ($SD = 1.04$, range = 1–5) and 1.58 ($SD = .81$, range = 1–5), respectively.

Tables 2, 3, and 4 present unadjusted and covariate-adjusted associations between predictors of health-related worries, functional impairment, and acute stress, respectively. In adjusted models, inability to obtain health care, lost wages, prior mental or physical health ailments, younger age, Hispanic ethnicity, and female gender were associated with acute stress, health-related worries, and functional impairment. Health care essential workers reported less functional impairment and acute stress compared with non-health care essential workers. Need for childcare was positively associated with acute stress symptoms, although not associated with health-related worries or functional impairment. Residents in the Northeast reported more health-related worries and acute stress than residents in the South. Those who completed the survey later reported more health-related worries, functional impairment, and acute stress compared with those who completed the survey in the earlier phases of the COVID-19 outbreak. Supplemental analyses indicated several statistically significant interaction terms. For functional impairment, need for childcare was moderated by essential worker status ($b = -.50$, 95% confidence interval, CI [–0.90, –0.11]). For health-related worries and acute stress, lost wages was moderated by essential worker status ($b =$

–.42, 95% CI [–0.77, –0.07]) and ($b = -0.32$, 95% CI [–0.56, –0.08]), respectively. See online supplemental data for full results of these supplemental analyses including tables (online Supplemental Materials Tables S1, S2, and S3) and figures (online Supplemental Materials Figures S1, S2, and S3).

Discussion

We explored predictors of psychosocial outcomes in health care and non-health care essential workers working in face-to-face interactions during the early phase of the COVID-19 outbreak in the United States. Results indicated secondary stressors (inability to obtain health care, lost wages), prior mental health ailments, prior physical-health diagnoses, and demographic indicators (younger age, female gender, and Hispanic ethnicity) were associated with greater functional impairment, health-related worries, and acute stress. Effect sizes were largest for difficulty obtaining health care, lost wages, age, female gender, and prior mental health ailments for acute stress; inability to obtain health care, and prior mental and physical health ailments for functional impairment; and inability to obtain health care, lost wages, age, prior mental and physical health ailments, and female gender for health-related worries. Contrary to our hypothesis, non-health care essential workers reported more functional impairment and acute stress than health care workers, yet effect sizes were relatively modest compared with other factors. Moreover, two stressors (needed

Table 2

Predictors of Health-Related Worries in Essential Workers During COVID-19 (N = 1,821)

Variable	Bivariate associations				Adjusted multivariate model			
	β	b	95% CI	p	β	b	95% CI	p
Health care worker ^a	0.01	0.03	[–0.14, 0.19]	.741	–0.01	–0.05	[–0.21, 0.12]	.564
Needed childcare	0.07	0.25	[0.09, 0.41]	.003	0.04	0.13	[–0.04, 0.29]	.126
Unable to obtain health care	0.16	0.56	[0.39, 0.73]	<.001	0.12	0.41	[0.25, 0.58]	<.001
Lost wages	0.11	0.29	[0.17, 0.40]	<.001	0.09	0.22	[0.10, 0.33]	<.001
Prior mental health dx ^b	0.14	0.38	[0.25, 0.50]	<.001	0.09	0.24	[0.12, 0.37]	<.001
Prior physical health dx ^c	0.04	0.04	[–0.004, 0.09]	.074	0.09	0.10	[0.05, 0.15]	<.001
Age	–0.10	–0.01	[–0.01, –0.004]	<.001	–0.09	–0.01	[–0.01, –0.003]	.001
Race/ethnicity ^d								
Black, non-Hispanic	0.02	0.08	[–0.08, 0.24]	.352	0.01	0.03	[–0.13, 0.20]	.673
Other/2+ races, non-Hispanic	0.06	0.24	[0.07, 0.42]	.007	0.04	0.15	[–0.02, 0.32]	.094
Hispanic	0.09	0.34	[0.17, 0.50]	<.001	0.08	0.29	[0.13, 0.46]	.001
Education ^e								
High school graduate	–0.16	–0.47	[–0.83, –0.11]	.011	–0.09	–0.25	[–0.60, 0.10]	.155
Some college	–0.25	–0.52	[–0.87, –0.18]	.003	–0.16	–0.33	[–0.67, 0.002]	.051
BA or above	–0.22	–0.48	[–0.83, –0.13]	.007	–0.12	–0.25	[–0.59, 0.09]	.154
Female gender	0.16	0.33	[0.24, 0.43]	<.001	0.14	0.29	[0.20, 0.39]	<.001
Income	–0.05	–0.03	[–0.06, –0.005]	.022	0.01	0.003	[–0.03, 0.03]	.828
Region ^f								
Midwest	–0.07	–0.16	[–0.31, –0.002]	.047	–0.06	–0.13	[–0.28, 0.02]	.085
South	–0.08	–0.18	[–0.34, –0.03]	.018	–0.09	–0.21	[–0.35, –0.06]	.006
West	–0.04	–0.10	[–0.26, 0.05]	.199	–0.06	–0.14	[–0.30, 0.01]	.070
Cohort ^g								
March 29, 2020 through April 7, 2020	0.11	0.25	[0.14, 0.37]	<.001	0.10	0.21	[0.10, 0.32]	<.001
April 8, 2020 through April 18, 2020	0.11	0.24	[0.12, 0.35]	<.001	0.09	0.19	[0.08, 0.30]	.001
Constant	—	—	—	—		2.67	[2.28, 3.08]	<.001
Model statistics							$F(20, 1,779) = 10.24, p < .001, R^2 = .10$	

Note. COVID-19 = coronavirus disease 2019; CI = confidence interval.

^a Non-health care essential worker = 0; ^b emotional/nervous/psychiatric problem, anxiety disorder, and depression (0 = none; 1 = one or more); ^c high cholesterol, hypertension, diabetes/high blood sugar, heart disease, stroke, cancer, lung disease, and other diagnoses (0–8); ^d White = 0; ^e <high school = 0; ^f Northeast = 0; ^g Cohort 1: March 18, 2020 through March 28, 2020 = 0.

Table 3
Predictors of Functional Impairment in Essential Workers During COVID-19 (N = 1,821)

Variable	Bivariate associations				Adjusted multivariate model			
	β	<i>b</i>	95% CI	<i>p</i>	β	<i>b</i>	95% CI	<i>p</i>
Health care worker ^a	-0.06	-0.17	[-0.29, -0.04]	.010	-0.06	-0.17	[-0.30, -0.04]	.009
Needed childcare	0.07	0.19	[0.06, 0.32]	.003	0.03	0.09	[-0.03, 0.22]	.145
Unable to obtain health care	0.21	0.59	[0.46, 0.71]	<.001	0.16	0.46	[0.33, 0.58]	<.001
Lost wages	0.09	0.19	[0.10, 0.29]	<.001	0.06	0.12	[0.04, 0.21]	.006
Prior mental health dx ^b	0.20	0.44	[0.34, 0.54]	<.001	0.16	0.35	[0.25, 0.45]	<.001
Prior physical health dx ^c	0.09	0.07	[0.03, 0.11]	<.001	0.12	0.10	[0.06, 0.14]	<.001
Age	-0.09	-0.01	[-0.01, -0.003]	<.001	-0.08	-0.01	[-0.01, -0.002]	.001
Race/ethnicity ^d								
Black, non-Hispanic	0.05	0.14	[0.01, 0.26]	.029	0.04	0.10	[-0.03, 0.22]	.133
Other/2+ races, non-Hispanic	0.09	0.27	[0.14, 0.41]	<.001	0.05	0.15	[0.02, 0.28]	.027
Hispanic	0.12	0.34	[0.22, 0.47]	<.001	0.10	0.28	[0.15, 0.41]	<.001
Education ^e								
High school graduate	-0.12	-0.27	[-0.55, 0.01]	.062	-0.05	-0.12	[-0.39, 0.15]	.391
Some college	-0.16	-0.26	[-0.54, 0.01]	.061	-0.08	-0.13	[-0.39, 0.13]	.339
BA or above	-0.17	-0.29	[-0.56, -0.01]	.041	-0.05	-0.09	[-0.36, 0.17]	.495
Female gender	0.09	0.16	[0.08, 0.23]	<.001	0.07	0.12	[0.04, 0.19]	.002
Income	-0.09	-0.04	[-0.07, -0.02]	<.001	-0.02	-0.01	[-0.03, 0.02]	.522
Region ^f								
Midwest	-0.02	-0.04	[-0.17, 0.08]	.479	-0.02	-0.04	[-0.15, 0.08]	.542
South	-0.01	-0.01	[-0.13, 0.11]	.839	-0.03	-0.05	[-0.17, 0.06]	.362
West	0.04	0.07	[-0.05, 0.19]	.251	0.01	0.01	[-0.11, 0.13]	.857
Cohort ^g								
March 29, 2020 through April 7, 2020	0.05	0.09	[-0.004, 0.18]	.063	0.03	0.06	[-0.03, 0.14]	.215
April 8, 2020 through April 18, 2020	0.07	0.13	[0.04, 0.22]	.005	0.05	0.09	[0.01, 0.18]	.034
Constant	—	—	—	—	—	1.60	[1.29, 1.92]	<.001
Model statistics	<i>F</i> (20, 1,776) = 12.64, <i>p</i> < .001, <i>R</i> ² = .12							

Note. COVID-19 = coronavirus disease 2019; CI = confidence interval.

^a Non-health care essential worker = 0; ^b emotional/nervous/psychiatric problem, anxiety disorder, and depression (0 = none; 1 = one or more); ^c high cholesterol, hypertension, diabetes/high blood sugar, heart disease, stroke, cancer, lung disease, and other diagnoses (0–8); ^d White = 0; ^e <high school = 0; ^f Northeast = 0; ^g Cohort March 18, 2020 through March 28, 2020 = 0.

childcare and lost wages) exhibited moderating effects, with negative outcomes exacerbated for non-health care workers.

Overall, distress and impairment were relatively low, indicating that while some reported distress and impairment, resilience early in the pandemic was common, in alignment with prior disaster research (Silver & Garfin, 2016). Indeed, people reported functional impairment, on average, between “not at all” and “a little of the time,” perhaps not surprising since these individuals were still working at the time of the assessment. Participants reported acute stress symptoms, on average, “just a little;” the mean for health related worries was highest of the variables assessed, with participants reporting these symptoms, on average, around the midpoint of the scale (i.e., between “rarely” and “sometimes”). Of note, the range and standard deviations indicate that despite resilience in many, some individuals reported distress and impairment at the scale maximums.

In a recent rapid review of research on health care workers employed during the COVID-19 pandemic, 24 studies identified risk factors for maladaptive psychological outcomes of health care workers—many of which paralleled the significant indicators presented herein, including female gender, younger age, and prior physical illness (De Kock et al., 2021). We expand this research by using data drawn from a probability-based sample of Americans that included both health care and non-health care essential workers. In the present study, female essential workers reported significantly higher amounts of all three psychosocial concerns compared with men, echoing previous studies conducted with COVID-19 health care workers (Tan et al., 2020) and during other

viral epidemics (Serrano-Ripoll et al., 2020). As in prior epidemics, history of mental health ailments was associated with negative outcomes (Asmundson et al., 2020; Elbay et al., 2020); however, present analyses highlight the additional role of preexisting physical health conditions on psychosocial responses. Age was negatively correlated with health-related worries, functional impairment, and acute stress, potentially due to fewer years of work experience and preparation, such as medical students entering the workforce early (Konda et al., 2020), nurses with junior titles (Steege et al., 2009), or other entry-level jobs (Elbay et al., 2020). Outcomes among all essential workers worsened as the pandemic advanced, suggesting psychosocial difficulties mirrored the outbreak’s U.S. progression.

Residents from the U.S. Northeast experienced more health-related worries and acute stress than Southern residents, aligning with regional outbreak severity during March and April 2020 when the data were collected (Konda et al., 2020). Hispanic ethnicity was positively associated with health-related worries, functional impairment, and acute stress. This aligns with data showing Hispanics, compared with other ethnicities, were more likely to live in a home where at least one worker was unable to work from home (Selden & Berdahl, 2020). Lack of childcare due to COVID-19 school closures was associated with higher acute stress symptoms, paralleling findings among a general sample that included health care workers that childcare concerns were associated with heightened anxiety and other worries (Hacimusalar et al., 2020). Notably, health care workers reported significantly *less* acute stress and functional impairment than non-health care

Table 4
Predictors of Acute Stress in Essential Workers During COVID-19 (N = 1,821)

Variable	Bivariate associations				Adjusted multivariate model			
	β	<i>b</i>	95% CI	<i>p</i>	β	<i>b</i>	95% CI	<i>P</i>
Health care worker ^a	-0.02	-0.05	[-0.16, 0.06]	.390	-0.06	-0.14	[-0.25, 0.27]	.015
Needed childcare	0.14	0.35	[0.24, 0.46]	<.001	0.09	0.22	[0.12, 0.33]	<.001
Unable to obtain health care	0.19	0.46	[0.35, 0.58]	<.001	0.13	0.33	[0.23, 0.44]	<.001
Lost wages	0.15	0.26	[0.18, 0.34]	<.001	0.11	0.20	[0.12, 0.28]	<.001
Prior mental health dx ^b	0.18	0.35	[0.27, 0.44]	<.001	0.12	0.24	[0.15, 0.33]	<.001
Prior physical health dx ^c	0.01	0.01	[-0.02, 0.04]	.678	0.08	0.06	[0.03, 0.09]	.001
Age	-0.17	-0.01	[-0.01, -0.01]	<.001	-0.14	-0.007	[-0.01, -0.005]	<.001
Race/ethnicity ^d								
Black, non-Hispanic	0.04	0.09	[-0.02, 0.20]	.121	0.02	0.04	[-0.07, 0.15]	.439
Other/2+ races, non-Hispanic	0.06	0.16	[0.04, 0.28]	.009	0.02	0.06	[-0.05, 0.18]	.297
Hispanic	0.10	0.24	[0.12, 0.35]	<.001	0.08	0.18	[0.07, 0.30]	.001
Education ^e								
High school graduate	-0.19	-0.38	[-0.63, -0.13]	.003	-0.10	-0.19	[-0.42, 0.04]	.106
Some college	-0.25	-0.37	[-0.61, -0.13]	.002	-0.14	-0.20	[-0.42, 0.02]	.081
BA or above	-0.21	-0.31	[-0.55, -0.07]	.011	-0.07	-0.10	[-0.33, 0.13]	.393
Female gender	0.18	0.26	[0.19, 0.32]	<.001	0.15	0.22	[0.16, 0.29]	<.001
Income	-0.05	-0.02	[-0.04, -0.002]	.029	0.02	0.01	[-0.01, 0.03]	.325
Region ^f								
Midwest	-0.06	-0.09	[-0.20, 0.01]	.085	-0.05	-0.09	[-0.19, 0.02]	.098
South	-0.06	-0.10	[-0.20, 0.01]	.067	-0.08	-0.12	[-0.22, -0.02]	.015
West	-0.02	-0.03	[-0.14, 0.08]	.633	-0.04	-0.06	[-0.16, 0.04]	.241
Cohort ^g								
March 29, 2020 through April 7, 2020	0.09	0.14	[0.06, 0.22]	.001	0.07	0.10	[0.03, 0.18]	.007
April 8, 2020 through April 18, 2020	0.10	0.16	[0.08, 0.24]	<.001	0.08	0.11	[0.05, 0.21]	.001
Constant						1.99	[1.72, 2.26]	<.001
Model statistics							$F(20, 1,782) = 15.90, p < .001, R^2 = .15$	

Note. COVID-19 = coronavirus disease 2019; CI = confidence interval.

^a Non-health care essential worker = 0; ^b emotional/nervous/psychiatric problem, anxiety disorder, and depression (0 = none; 1 = one or more); ^c high cholesterol, hypertension, diabetes/high blood sugar, heart disease, stroke, cancer, lung disease, and other diagnoses (0–8); ^d White = 0; ^e <high school = 0; ^f Northeast = 0; ^g Cohort March 18, 2020 through March 28, 2020 = 0.

essential workers, perhaps because health care workers have more protocols for working with infectious diseases, greater meaning in their work, or more experience on the job. Indeed, physicians in Izmir/Turkey who were actively treating COVID-19 had lower burnout and higher feelings of accomplishments compared with physicians treating other conditions (Dinibutun, 2020). Moreover, adequate protection training for nurses in China was associated with lower anxiety, acute stress, and depression in nurses during early phases of COVID-19 (Cai et al., 2020). These factors may be more likely explanations than job experience, as burnout and stress often increases in health care workers over their career trajectory (Cull et al., 2019).

Strengths and Limitations

Key strengths of this study include a large sample of U.S. essential workers, derived from an overall representative group of Americans recruited during COVID-19's early progression in the United States. Given the dearth of data on non-health care essential workers actively employed during COVID-19, our contributions to the literature demonstrate that this group may be critical to target with mental health resources and outreach efforts. This is particularly crucial as society enters a new phase of the COVID-19 pandemic, where emerging variants are more contagious and community transmission remains high (Christie et al., 2021), yet essential workers are still required to engage in face-to-face interactions, potentially with fewer protections (e.g., relaxed mask mandates). Our analyses

included prepandemic physical and mental health diagnosis data, allowing us to draw comparisons between prior diagnoses and present psychological symptomatology without the bias inherent in retrospective reporting.

We acknowledge several limitations. Our data on exposure and symptoms were collected concurrently. While the overall sample of Americans we surveyed was representative of the United States and the subsample of essential workers we surveyed is likely proportional to their representation in the general population, we cannot consider this a representative sample of essential workers. While physical and mental health data were collected in advance of the pandemic, physical health data was available as count of potential ailments and mental health was available as a dichotomous yes/no variable. Finally, we did not assess specific job, seniority, or workplace safety precautions, which may further explain variability in outcomes.

Conclusions and Future Directions

To our knowledge, this is the first study evaluating psychosocial outcomes in both health care and non-health care U.S. essential workers during the COVID-19 pandemic using a probability-based design. While many workplaces have implemented COVID-19 mitigation strategies, health and safety concerns remain, and may be associated with psychosocial difficulties. Indeed, acute psychosocial responses to stress may portend deleterious downstream mental and physical health effects, including depression, anxiety, and cardiovascular problems, as well as reduced working capacity and lower self-

reported overall health (Garfin et al., 2018). COVID-19-related psychological and occupational impairment could be eased by implementing reliable public health workplace policies (Tan et al., 2020) that encourage healthy coping strategies. Some hospitals have implemented innovative mental health and resilience-building centers for health care workers in response to the pandemic (DePierro et al., 2020), which non-health care sectors should consider enacting. Workers should be provided adequate benefits like health insurance and paid sick leave. The COVID-19 pandemic also ushered in advances in telehealth and online resources to improve self-care; these include individual and group psychotherapy and wellness apps that can be utilized on mobile devices (Garfin, 2020). Such resources may be valuable for workers as they allow for the safe delivery of interventions and greater convenience for scheduling around work hours. Attention should be paid to non-health care essential workers in addition to health care essential workers, as their likelihood for functional impairment may be heightened without corresponding supports. As COVID-19 becomes endemic, appropriate resources should be provided for workers who are most vulnerable to COVID-19 and as we strive to prepare for future outbreaks.

References

- Asmundson, G. J. G., Paluszek, M. M., Landry, C. A., Rachor, G. S., McKay, D., & Taylor, S. (2020). Do pre-existing anxiety-related and mood disorders differentially impact COVID-19 stress responses and coping? *Journal of Anxiety Disorders, 74*, 102271. <https://doi.org/10.1016/j.janxdis.2020.102271>
- Baker, M. G., Peckham, T. K., & Seixas, N. S. (2020). Estimating the burden of United States workers exposed to infection or disease: A key factor in containing risk of COVID-19 infection. *PLoS ONE, 15*(4), e0232452. <https://doi.org/10.1371/journal.pone.0232452>
- Benfante, A., Di Tella, M., Romeo, A., & Castelli, L. (2020). Traumatic stress in healthcare workers during COVID-19 pandemic: A review of the immediate impact. *Frontiers in Psychology, 11*, 569935. <https://doi.org/10.3389/fpsyg.2020.569935>
- Bryant, R. A. (2016). *Acute stress disorder: What it is and how to treat it*. New York: Guilford.
- Cabarkapa, S., Nadjidai, S. E., Murgier, J., & Ng, C. H. (2020). The psychological impact of COVID-19 and other viral epidemics on frontline healthcare workers and ways to address it: A rapid systematic review. *Brain, Behavior, and Immunity, 88*, 100144. <https://doi.org/10.1016/j.bbih.2020.100144>
- Cai, Z., Cui, Q., Liu, Z., Li, J., Gong, X., Liu, J., Wan, Z., Yuan, X., Li, X., Chen, C., & Wang, G. (2020). Nurses endured high risks of psychological problems under the epidemic of COVID-19 in a longitudinal study in Wuhan China. *Journal of Psychiatric Research, 131*, 132–137. <https://doi.org/10.1016/j.jpsychires.2020.09.007>
- Chew, N. W. S., Lee, G. K. H., Tan, B. Y. Q., Jing, M., Goh, Y., Ngiam, N. J. H., Yeo, L. L. L., Ahmad, A., Ahmed Khan, F., Napoleon Shanmugam, G., Sharma, A. K., Komalkumar, R. N., Meenakshi, P. V., Shah, K., Patel, B., Chan, B. P. L., Sunny, S., Chandra, B., Ong, J. J. Y., . . . Sharma, V. K. (2020). A multinational, multicentre study on the psychological outcomes and associated physical symptoms amongst healthcare workers during COVID-19 outbreak. *Brain, Behavior, and Immunity, 88*, 559–565. <https://doi.org/10.1016/j.bbi.2020.04.049>
- Chew, N. W. S., Ngiam, J. N., Tan, B. Y.-Q., Tham, S.-M., Tan, C. Y.-S., Jing, M., Sagayanathan, R., Chen, J. T., Wong, L. Y. H., Ahmad, A., Khan, F. A., Marmin, M., Hassan, F. B., Sharon, T. M. L., Lim, C. H., Mohaini, M. I. B., Danuaji, R., Nguyen, T. H., Tsvigoulis, G., . . . Sharma, V. K. (2020). Asian-Pacific perspective on the psychological well-being of healthcare workers during the evolution of the COVID-19 pandemic. *BJPsych Open, 6*(6), e116. <https://doi.org/10.1192/bjo.2020.98>
- Christie, A., Brooks, J. T., Hicks, L. A., Sauber-Schatz, E. K., Yoder, J. S., & Honein, M. A., & The CDC COVID-19 Response Team. (2021). Guidance for implementing COVID-19 prevention strategies in the context of varying community transmission levels and vaccination coverage. *Morbidity and Mortality Weekly Report, 70*(30), 1044–1047. <https://doi.org/10.15585/mmwr.mm7030e2>
- Cull, W. L., Frintner, M. P., Starmer, A. J., & Leslie, L. K. (2019). Longitudinal analyses of pediatrician burnout. *Academic Pediatrics, 19*(3), 256–262. <https://doi.org/10.1016/j.acap.2018.11.006>
- De Kock, J. H., Latham, H. A., Leslie, S. J., Grindle, M., Munoz, S.-A., Ellis, L., Polson, R., & O'Malley, C. M. (2021). A rapid review of the impact of COVID-19 on the mental health of healthcare workers: Implications for supporting psychological well-being. *BMC Public Health, 21*(1), 104. <https://doi.org/10.1186/s12889-020-10070-3>
- DePierro, J., Katz, C. L., Marin, D., Feder, A., Bevilacqua, L., Sharma, V., Hurtado, A., Ripp, J., Lim, S., & Charney, D. (2020). Mount Sinai's Center for Stress, Resilience and Personal Growth as a model for responding to the impact of COVID-19 on health care workers. *Psychiatry Research, 293*, 113426. <https://doi.org/10.1016/j.psychres.2020.113426>
- Dinibutun, S. R. (2020). Factors associated with burnout among physicians: an evaluation during a period of COVID-19 pandemic. *Journal of Healthcare Leadership, 12*, 85–94. <https://doi.org/10.2147/JHL.S270440>
- Elbay, R. Y., Kurtulmuş, A., Arpacıoğlu, S., & Karadere, E. (2020). Depression, anxiety, stress levels of physicians and associated factors in Covid-19 pandemics. *Psychiatry Research, 290*, 113130. <https://doi.org/10.1016/j.psychres.2020.113130>
- Flaherty, G. T., Hession, P., Liew, C. H., Lim, B. C. W., Leong, T. K., Lim, V., & Sulaiman, L. H. (2020). COVID-19 in adult patients with pre-existing chronic cardiac, respiratory and metabolic disease: A critical literature review with clinical recommendations. *Tropical Diseases, Travel Medicine and Vaccines, 6*(1), 16. <https://doi.org/10.1186/s40794-020-00118-y>
- Galea, S., Brewin, C. R., Gruber, M., Jones, R. T., King, D. W., King, L. A., McNally, R. J., Ursano, R. J., Petukhova, M., & Kessler, R. C. (2007). Exposure to hurricane-related stressors and mental illness after Hurricane Katrina. *Archives of General Psychiatry, 64*(12), 1427–1434. <https://doi.org/10.1001/archpsyc.64.12.1427>
- Gallagher, S., Roy, A., Domeracki, S. J., Mohrmann, T., Missar, V., Jule, J., Sharma, S., & DeWitt, R. (2021). The low-wage essential worker: Occupational concerns and needs in the COVID-19 pandemic—A round table. *Workplace Health & Safety, 69*(4), 154–160. <https://doi.org/10.1177/2165079920988682>
- Garfin, D. R. (2020). Technology as a coping tool during the coronavirus disease 2019 (COVID-19) pandemic: Implications and recommendations. *Stress and Health, 36*(4), 555–559. <https://doi.org/10.1002/smi.2975>
- Garfin, D. R., Silver, R. C., Ugalde, F. J., Linn, H., & Inostroza, M. (2014). Exposure to rapid succession disasters: A study of residents at the epicenter of the Chilean Bío Bío earthquake. *Journal of Abnormal Psychology, 123*(3), 545–556. <https://doi.org/10.1037/a0037374>
- Garfin, D. R., Thompson, R. R., & Holman, E. A. (2018). Acute stress and subsequent health outcomes: A systematic review. *Journal of Psychosomatic Research, 112*, 107–113. <https://doi.org/10.1016/j.jpsychores.2018.05.017>
- Ghilarducci, T., & Farmand, A. (2020). Older workers on the COVID-19 frontlines without paid sick leave. *Journal of Aging & Social Policy, 32*(4–5), 471–476. <https://doi.org/10.1080/08959420.2020.1765685>
- Grover, S., Mehra, A., Sahoo, S., Avasthi, A., Rao, T. S. S., Vaishnav, M., Dalal, P. K., Saha, G., Singh, O. P., Chakraborty, K., Reedy, Y. C. J., Rao, N. P., Tripathi, A., Chadda, R. K., Mishra, K. K., Rao, G. P., Kumar, V., Gautam, S., Sarkar, S., . . . & Subramanyam, A. (2021). Evaluation of psychological impact of COVID-19 on health-care workers. *Indian Journal of Psychiatry, 63*(3), 222–227. https://doi.org/10.4103/indianjpsychiatry.indianjpsychiatry_1129_20

- Hacimusalar, Y., Kahve, A. C., Yasar, A. B., & Aydin, M. S. (2020). Anxiety and hopelessness levels in COVID-19 pandemic: A comparative study of healthcare professionals and other community sample in Turkey. *Journal of Psychiatric Research, 129*, 181–188. <https://doi.org/10.1016/j.jpsychires.2020.07.024>
- Holman, E. A., Thompson, R. R., Garfin, D. R., & Silver, R. C. (2020). The unfolding COVID-19 pandemic: A probability-based, nationally representative study of mental health in the United States. *Science Advances, 6*(42), eabd5390. <https://doi.org/10.1126/sciadv.abd5390>
- Kantamneni, N. (2020). The impact of the COVID-19 pandemic on marginalized populations in the United States: A research agenda. *Journal of Vocational Behavior, 119*, 103439. <https://doi.org/10.1016/j.jvb.2020.103439>
- Kessler, R. C., McLaughlin, K. A., Koenen, K. C., Petukhova, M., & Hill, E. D., & The WHO World Mental Health Survey Consortium. (2012). The importance of secondary trauma exposure for post-disaster mental disorder. *Epidemiology and Psychiatric Sciences, 21*(1), 35–45. <https://doi.org/10.1017/S2045796011000758>
- Kofman, A., Kantor, R., & Adashi, E. Y. (2021). Potential COVID-19 endgame scenarios eradication, elimination, cohabitation, or conflagration? *Journal of the American Medical Association, 326*, 303–304. <https://doi.org/10.1001/jama.2021.11042>
- Konda, S. R., Dankert, J. F., Merkow, D., Lin, C. C., Kaplan, D. J., Haskel, J. D., & Ganta, A. (2020). COVID-19 response in the global epicenter: Converting a New York City level 1 orthopedic trauma service into a hybrid orthopedic and medicine COVID-19 management team. *Journal of Orthopaedic Trauma, 34*(8), 411–417. <https://doi.org/10.1097/BOT.0000000000001792>
- Lai, J., Ma, S., Wang, Y., Cai, Z., Hu, J., Wei, N., Wu, J., Du, H., Chen, T., Li, R., Tan, H., Kang, L., Yao, L., Huang, M., Wang, H., Wang, G., Liu, Z., & Hu, S. (2020). Factors associated with mental health outcomes among health care workers exposed to Coronavirus Disease 2019. *JAMA Network Open, 3*(3), e203976. <https://doi.org/10.1001/jamanetworkopen.2020.3976>
- Luo, Y., Chua, C. R., Xiong, Z., Ho, R. C., & Ho, C. S. H. (2020). A Systematic review of the impact of viral respiratory epidemics on mental health: An implication on the coronavirus disease 2019 pandemic. *Frontiers in Psychiatry, 11*, 565098. <https://doi.org/10.3389/fpsy.2020.565098>
- Magill, E., Siegel, Z., & Pike, K. (2020). The mental health of frontline healthcare providers during pandemics: A rapid review of literature. *Psychiatric Services, 71*(12), 1260–1269. <https://doi.org/10.1017/CBO9781107415324.004>
- McGinty, E. E., Presskreischer, R., Anderson, K. E., Han, H., & Barry, C. L. (2020). Psychological distress and covid-19-related stressors reported in a longitudinal cohort of U.S. adults in April and July 2020. *Journal of the American Medical Association, 324*(24), 2555–2557. <https://doi.org/10.1001/jama.2020.21231>
- Pappa, S., Ntella, V., Giannakas, T., Giannakoulis, V. G., Papoutsis, E., & Katsaounou, P. (2020). Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain, Behavior, and Immunity, 88*, 901–907. <https://doi.org/10.1016/j.bbi.2020.05.026>
- Rogalski, M. A., Gowler, C. D., Shaw, C. L., Hufbauer, R. A., & Duffy, M. A. (2017). Human drivers of ecological and evolutionary dynamics in emerging and disappearing infectious disease systems. *Philosophical Transactions of the Royal Society B: Biological Sciences, 372*, 20160043. <https://doi.org/10.1098/rstb.2016.0043>
- Ruiz-Frutos, C., Ortega-Moreno, M., Allande-Cussó, R., Domínguez-Salas, S., Dias, A., & Gómez-Salgado, J. (2021). Health-related factors of psychological distress during the COVID-19 pandemic among non-health workers in Spain. *Safety Science, 133*, 104996. <https://doi.org/10.1016/j.ssci.2020.104996>
- Sallam, M. (2021). Covid-19 vaccine hesitancy worldwide: A concise systematic review of vaccine acceptance rates. *Vaccines, 9*(2), 160. <https://doi.org/10.3390/vaccines9020160>
- Selden, T. M., & Berdahl, T. A. (2020). COVID-19 and racial/ethnic disparities in health risk, employment, and household composition. *Health Affairs, 39*(9), 1624–1632. <https://doi.org/10.1377/hlthaff.2020.00897>
- Serrano-Ripoll, M. J., Meneses-Echavez, J. F., Ricci-Cabello, I., Fraile-Navarro, D., Fiol-deRoque, M. A., Pastor-Moreno, G., Castro, A., Ruiz-Pérez, I., Zamanillo Campos, R., & Gonçalves-Bradley, D. C. (2020). Impact of viral epidemic outbreaks on mental health of healthcare workers: A rapid systematic review and meta-analysis. *Journal of Affective Disorders, 277*, 347–357. <https://doi.org/10.1016/j.jad.2020.08.034>
- Shanafelt, T., Ripp, J., & Trockel, M. (2020). Understanding and addressing sources of anxiety among health care professionals during the COVID-19 pandemic. *Journal of the American Medical Association, 323*, 2133–2134. <https://doi.org/10.1001/jama.2020.5893>
- Silver, R. C., & Garfin, D. R. (2016). Coping with disasters. In J. C. Norcross, G. R. VandenBos, D. K. Freedheim, & N. Pole (Eds.), *APA handbook of clinical psychology: Psychopathology and health* (pp. 597–611). American Psychological Association. <https://doi.org/10.1037/14862-029>
- Sim, M. R. (2020). The COVID-19 pandemic: Major risks to healthcare and other workers on the front line. *Occupational and Environmental Medicine, 77*(5), 281–282. <https://doi.org/10.1136/oemed-2020-106567>
- Steege, A. L., Baron, S., Davis, S., Torres-Kilgore, J., & Sweeney, M. H. (2009). Pandemic influenza and farmworkers: The effects of employment, social, and economic factors. *American Journal of Public Health, 99*(2), S308–S315. <https://doi.org/10.2105/AJPH.2009.161091>
- Tan, W., Hao, F., McIntyre, R. S., Jiang, L., Jiang, X., Zhang, L., Zhao, X., Zou, Y., Hu, Y., Luo, X., Zhang, Z., Lai, A., Ho, R., Tran, B., Ho, C., & Tam, W. (2020). Is returning to work during the COVID-19 pandemic stressful? A study on immediate mental health status and psychoneuroimmunity prevention measures of Chinese workforce. *Brain, Behavior, and Immunity, 87*, 84–92. <https://doi.org/10.1016/j.bbi.2020.04.055>
- Tan, Y. Q., Wang, Z., Yap, Q. V., Chan, Y. H., Ho, R. C., Hamid, A. R. A. H., Garcia, B. N., & Chiong, E. (2021). Psychological health of surgeons in a time of COVID-19. *Annals of Surgery*. Advance online publication. <https://doi.org/10.1097/sla.0000000000004775>
- Thompson, R. R., Garfin, D. R., Holman, E. A., & Silver, R. C. (2017). Distress, worry, and functioning following a global health crisis: A national study of Americans' responses to Ebola. *Clinical Psychological Science, 5*(3), 513–521. <https://doi.org/10.1177/2167702617692030>
- Tian, F., Li, H., Tian, S., Yang, J., Shao, J., & Tian, C. (2020). Psychological symptoms of ordinary Chinese citizens based on SCL-90 during the level I emergency response to COVID-19. *Psychiatry Research, 288*, 112992. <https://doi.org/10.1016/j.psychres.2020.112992>
- Toh, W. L., Meyer, D., Phillipou, A., Tan, E. J., Van Rheenen, T. E., Neill, E., & Rossell, S. L. (2021). Mental health status of healthcare versus other essential workers in Australia amidst the COVID-19 pandemic: Initial results from the collate project. *Psychiatry Research, 298*, 113822. <https://doi.org/10.1016/j.psychres.2021.113822>
- Ware, J. E., & Sherbourne, C. D. (1992). The MOS 36-Item Short-Form Health Survey (SF-36): I. Conceptual frame-work and item selection. *Medical Care, 30*, 473–483. <https://doi.org/10.1097/00005650-199206000-00002>
- Wright, T. A., & Cropanzano, R. (2000). Psychological well-being and job satisfaction as predictors of job performance. *Journal of Occupational Health Psychology, 5*(1), 84–94. <https://doi.org/10.1037/1076-8998.5.1.84>
- Zheng, J., Morstead, T., Sin, N., Klaiber, P., Umberson, D., Kamble, S., & DeLongis, A. (2021). Psychological distress in North America during COVID-19: The role of pandemic-related stressors. *Social Science and Medicine, 270*, 113687. <https://doi.org/10.1016/j.socscimed.2021.113687>

Received August 12, 2021

Revision received October 30, 2021

Accepted November 16, 2021 ■